

Poisson algebras associated to quasi-Hopf algebras

Benjamin Enriquez and Gilles Halbout

Institut de Recherche Mathématique Avancée de Strasbourg
 UMR 7501 de l'Université Louis Pasteur et du CNRS
 7, rue R. Descartes F-67084 Strasbourg
 e-mail: enriquez@math.u-strasbg.fr
 e-mail: halbout@math.u-strasbg.fr

February 1, 2008

Abstract

We define admissible quasi-Hopf quantized universal enveloping (QHQUE) algebras by $\hbar$ -adic valuation conditions. We show that any QHQUE algebra is twist-equivalent to an admissible one. We prove a related statement: any associator is twist-equivalent to a Lie associator. We attach a quantized formal series algebra to each admissible QHQUE algebra and study the resulting Poisson algebras.

§ 0 Introduction

In [WX], Weinstein and Xu introduced a geometric counterpart of quasitriangular quantum groups: they proved that if $(\mathfrak{g}, r)$ is a finite dimensional quasi-triangular Lie bialgebra, then the dual group G^* is equipped with a braiding $\mathcal{R}_{WX}$ with properties analogous to those of quantum R -matrices (in particular, it is a set-theoretic solution of the quantum Yang-Baxter Equation). An explicit relation to the theory of quantum groups was later given in [GH, EH, EGH]: to a quasi-triangular QUE algebra $(U_{\hbar}(\mathfrak{g}), m, R)$ quantizing $(\mathfrak{g}, r)$, one associates its quantized formal series algebra (QFSA) $U_{\hbar}(\mathfrak{g})' \subset U_{\hbar}(\mathfrak{g})$; $U_{\hbar}(\mathfrak{g})'$ is a flat deformation of the Hopf-co-Poisson algebra $\mathcal{O}_{G^*} = (U(\mathfrak{g}^*))^*$ of formal functions of G^* . Then one proves that $\text{Ad}(R)$ preserves $U_{\hbar}(\mathfrak{g})'^{\otimes 2}$, and $\text{Ad}(R)|_{\hbar=0}$ coincides with the automorphism $\mathcal{R}_{WX}$ of $\mathcal{O}_{G^*}^{\otimes 2}$; moreover, $\rho = \hbar \log(R)|_{\hbar=0}$ is a function of $\mathcal{O}_{G^*}^{\otimes 2}$, independent on a quantization of $\mathfrak{g}^*$, which may be expressed universally in terms of r , and $\mathcal{R}_{WX}$ coincides with the “time one automorphism” of the Hamiltonian vector field generated by ρ .

In this paper, we study the analogous problem in the case of quasi-quantum groups (quasi-Hopf QUE algebras). The classical limit of a QHQUE algebra is a Lie quasibialgebra (LQBA). V. Drinfeld proposed to attach Poisson-Lie “quasi-groups” to each LQBA ([Dr4]). Axioms for Poisson-Lie quasi-groups are the quasi-Hopf analogues of the Weinstein-Xu axioms.

A *Poisson-Lie quasi-group* is a Poisson manifold X , together with a “product” Poisson map $X^2 \xrightarrow{m_X} X$, a unit for this product $e \in X$, and Poisson automorphisms $\Phi_X \in \text{Aut}(X^3)$,

$\Phi_X^{12,3,4}$, $\Phi_X^{1,23,4}$ and $\Phi_X^{1,2,34} \in \text{Aut}(X^4)$, such that

$$\begin{aligned} m_X \circ (\text{id} \times m_X) &= m_X \circ (m_X \times \text{id}) \circ \Phi_X, \\ (m_X \times \text{id} \times \text{id}) \circ \Phi_X^{12,3,4} &= \Phi_X \circ (m_X \times \text{id} \times \text{id}), \\ (\text{id} \times m_X \times \text{id}) \circ \Phi_X^{1,23,4} &= \Phi_X \circ (\text{id} \times m_X \times \text{id}), \text{ etc.} \\ \text{and } \Phi_X^{1,2,34} \circ \Phi_X^{12,3,4} &= (\text{id} \times \Phi_X) \circ \Phi_X^{1,23,4} \circ (\Phi_X \times \text{id}). \end{aligned}$$

A *twistor* for the quasi-group (X, m_X, Φ_X) is a collection of Poisson automorphisms $F_X \in \text{Aut}(X^2)$, $F_X^{12,3}$, $F_X^{1,23} \in \text{Aut}(X^3)$, $F_X^{(12)3,4}$, $F_X^{1(23),4}$, $F_X^{12,34}$, $F_X^{1(23),4}$, $F_X^{1,(23)4} \in \text{Aut}(X^4)$ such that

$$\begin{aligned} (m_X \times \text{id}) \circ F_X^{12,3} &= F_X \circ (m_X \times \text{id}), \\ ((m_X \circ (\text{id} \times m_X)) \times \text{id}) \circ F_X^{1(23),4} &= F_X \circ ((m_X \circ (\text{id} \times m_X)) \times \text{id}), \\ F_X^{(12)3,4} &= (\Phi_X \times \text{id}) \circ F_X^{1(23),4} \circ (\Phi_X \times \text{id})^{-1}, \text{ etc.} \end{aligned}$$

A twistor replaces the quasi-group (X, m_X, Φ_X) by (X, m'_X, Φ'_X) with $m'_X = m_X \circ F_X$ and $\Phi'_X = (F_X^{1,23})^{-1} \circ (F_X \times \text{id})^{-1} \circ \Phi_X \circ F_X^{1,23} \circ (\text{id} \times F_X)$.

(Other axioms for Poisson-Lie quasi-groups were proposed in a differential-geometric language in [Ban, KS].)

We do not know a “geometric” construction of a twist-equivalence class of (X, m_X, Φ_X) associated to each Lie quasi-bialgebra, in the spirit of [WX]. Instead we generalize the “construction of a QFS algebra and passage to Poisson geometry” part of the above discussion, and we derive from there a construction of triples (X, m_X, Φ_X) , in the case of Lie quasi-bialgebras with vanishing cobracket.

Let us describe the generalization of the “construction of a QFS algebra” part (precise statements are in Section 1). We introduce the notion of an *admissible* quasi-Hopf QUE algebra, and we associate a QFSA to such a QHQUE algebra. Each QHQUE algebra can be made admissible after a suitable twist.

We generalize the “passage to Poisson geometry” part as follows. The reduction modulo $\hbar$ of the obtained QFS algebra is a quintuple $(A, m, P, \Delta, \bar{\varphi})$ satisfying certain axioms; in particular $\exp(V_{\bar{\varphi}})$ is an automorphism of $A^{\otimes 3}$, and $(A, m, \exp(V_{\bar{\varphi}}))$ satisfies the axioms dual to those of (X, m_X, Φ_X) .

When the Lie quasi-bialgebra arises from a metrized Lie algebra, admissible QHQUE algebras quantizing it are given by Lie associators, and we obtain a quasi-group (X, m_X, Φ_X) using our construction. We also prove that its twist-equivalence class does not depend on the choice of an associator.

Finally, we prove a related result: any associator is twist-equivalent to a unique Lie associator.

§ 1 Outline of results

Let $\mathbb{K}$ be a field of characteristic 0. Let (U, m) be a topologically free $\mathbb{K}[[\hbar]]$ -algebra equipped with algebra morphisms

$$\Delta : U \rightarrow U \hat{\otimes} U, \text{ and } \varepsilon : U \rightarrow \mathbb{K}[[\hbar]]$$

$$\text{with } (\varepsilon \otimes \text{id}) \circ \Delta = (\text{id} \otimes \varepsilon) \circ \Delta = \text{id}$$

such that the reduction of (U, m, Δ) modulo $\hbar$ is a universal enveloping algebra. Set

$$U' = \{x \in U \mid \text{for any tree } P, \delta^{(P)}(x) \in \hbar^{|P|} U^{\otimes |P|}\}$$

(see the definitions of a tree, $\delta^{(P)}$, and $|P|$ in Section 2). We prove:

Theorem 1.1. *U' is a topologically free $\mathbb{K}[[\hbar]]$ -algebra. It is equipped with a complete decreasing algebra filtration*

$$(U')^{(n)} = \{x \in U' \mid \text{for any tree } P, \delta^{(P)}(x) \in \hbar^n U^{\otimes |P|}\}.$$

U' is stable under the multiplication m and the map $\Delta : U \rightarrow U^{\widehat{\otimes} 2}$ induces a continuous algebra morphism

$$\Delta_{U'} : U' \rightarrow U'^{\widehat{\otimes} 2} = \varprojlim_n \left(U'^{\widehat{\otimes} 2} / \sum_{p,q \mid p+q=n} U'^{(p)} \otimes U'^{(q)} \right).$$

Set $\mathcal{O} := U' / \hbar U'$. Then $\mathcal{O}$ is a complete commutative local ring and the reduction modulo $\hbar$ of $\Delta_{U'}$ is a continuous ring morphism

$$\Delta_{\mathcal{O}} : \mathcal{O} \rightarrow \mathcal{O}^{\widehat{\otimes} 2} = \varprojlim_n \left(\mathcal{O}^{\widehat{\otimes} 2} / \sum_{p,q \mid p+q=n} \mathcal{O}^{(p)} \otimes \mathcal{O}^{(q)} \right),$$

where $\mathcal{O}^{(p)} = U'^{(p)} / (\hbar U' \cap U'^{(p)})$.

Theorem 1.2. *Let (U, m, Δ, Φ) be a quasi-Hopf QUE algebra. Assume that*

$$\hbar \log(\Phi) \in (U')^{\widehat{\otimes} 3}. \quad (1.1)$$

Then there is a noncanonical isomorphism of filtered algebras $U' / \hbar U' \rightarrow \widehat{S}(\mathfrak{g})$, where $\widehat{S}(\mathfrak{g})$ is the formal series completion of the symmetric algebra $S(\mathfrak{g})$.

When (U, m, Δ, Φ) satisfies the hypothesis (1.1), we say that it is *admissible*. In that case, we say that U' is the quantized formal series algebra (QFSA) corresponding to (U, m, Δ, Φ) . Let us recall the notion of a *twist* of a quasi-Hopf QUE algebra (U, m, Δ, Φ) . This is an element $F \in (U^{\widehat{\otimes} 2})^\times$, such that $(\varepsilon \otimes \text{id})(F) = (\text{id} \otimes \varepsilon)(F) = 1$. It transforms (U, m, Δ, Φ) into the quasi-Hopf algebra $(U, m, {}^F \Delta, {}^F \Phi)$, where

$${}^F \Delta = \text{Ad}(F) \circ \Delta, \text{ and } {}^F \Phi = (1 \otimes F)(\text{id} \otimes \Delta)(F) \Phi (\Delta \otimes \text{id})(F)^{-1} (F \otimes 1)^{-1}.$$

Theorem 1.3.

- 1) *Let (U, m, Δ, Φ) be an admissible quasi-Hopf QUE algebra. Let us say that a twist F of U is admissible if $\hbar \log(F) \in U'^{\widehat{\otimes} 2}$. Then the twisted quasi-Hopf algebra $(U, m, {}^F \Delta, {}^F \Phi)$ is also admissible, and its QFSA coincides with U' .*
- 2) *Let (U, m, Δ, Φ) be an arbitrary quasi-Hopf QUE algebra. There exists a twist F_0 of U such that the twisted quasi-Hopf algebra $(U, m, {}^{F_0} \Delta, {}^{F_0} \Phi)$ is admissible.*

Theorem 1.3 can be interpreted as follows. Let (U, m) be a formal deformation of a universal enveloping algebra. The set of twists of U is a subgroup $\mathcal{T}$ of $(U^{\widehat{\otimes} 2})^\times$. Denote by $\mathcal{Q}$ the set of all quasi-Hopf structures on (U, m) , and by $\mathcal{Q}_{\text{adm}}$ the subset of admissible structures. If $\mathcal{Q}$ is nonempty, then $\mathcal{Q}_{\text{adm}}$ is also nonempty, and all its elements give rise

to the same subalgebra $U' \subset U$ (Theorem 1.3, 1)). Using U' , we then define the subgroup $\mathcal{T}_{\text{adm}} \subset \mathcal{T}$ of admissible twists. We have a natural action of $\mathcal{T}$ on $\mathcal{Q}$, which restricts to an action of $\mathcal{T}_{\text{adm}}$ on $\mathcal{Q}_{\text{adm}}$. Theorem 1.3 2) says that the natural map

$$\mathcal{Q}_{\text{adm}}/\mathcal{T}_{\text{adm}} \rightarrow \mathcal{Q}/\mathcal{T}$$

is surjective. Let us explain why it is not injective in general. Any QUE Hopf algebra (U, m, Δ) is admissible as a quasi-Hopf algebra. If $u \in U^\times$ and $F = (u \otimes u)\Delta(u)^{-1}$, then $(U, m, {}^F\Delta)$ is a Hopf algebra. So (U, m, Δ) and $(U, m, {}^F\Delta)$ are in the same class of $\mathcal{Q}/\mathcal{T}$. These are also two elements of $\mathcal{Q}_{\text{adm}}$; the corresponding QFS algebras are U' and $\text{Ad}(u)(U')$. In general, these algebras do not coincide, so (U, m, Δ) and $(U, m, {}^F\Delta)$ are not in the same class of $\mathcal{Q}_{\text{adm}}/\mathcal{T}_{\text{adm}}$.

The following result is a refinement of Proposition 3.10 of [Dr2]. Let $(\mathfrak{g}, \mu, \varphi)$ be a pair of a Lie algebra $(\mathfrak{g}, \mu)$ and $\varphi \in \wedge^3(\mathfrak{g})^\mathfrak{g}$. Then $(\mathfrak{g}, \delta = 0, \varphi)$ is a Lie bialgebra.

Proposition 1.4. *There exists a series $\mathcal{E}(\varphi) \in U(\mathfrak{g})^{\otimes 3}[[\hbar]]$, expressed in terms of (μ, φ) by universal acyclic expressions, such that $(U(\mathfrak{g})[[\hbar]], m_0, \Delta_0, \mathcal{E}(\varphi))$ is an admissible quantization of $(\mathfrak{g}, \mu, \varphi)$.*

This proposition is proved in Section 6.

Let us define a Drinfeld algebra as follows:

Definition 1.5. *A Drinfeld algebra is a quintuple $(A, m_0, P, \Delta, \tilde{\varphi})$, where*

- (A, m_0) is a formal series algebra,
- P is a Poisson structure on A “vanishing at the origin” (i.e., such that $\text{Im}(P) \subset \mathfrak{m}_A$, where $\mathfrak{m}_A$ is the maximal ideal of A),
- $\Delta : A \rightarrow A \hat{\otimes} A$ is a continuous Poisson algebra morphism, such that $(\varepsilon \otimes \text{id}) \circ \Delta = (\text{id} \otimes \varepsilon) \circ \Delta = \text{id}$, where $\varepsilon : A \rightarrow A/\mathfrak{m}_A = \mathbb{K}$ is the natural projection,
- $\tilde{\varphi} \in (\mathfrak{m}_A)^{\hat{\otimes} 3}$ satisfies

$$\begin{aligned} (\text{id} \otimes \Delta)(\Delta(a)) &= \tilde{\varphi} \star (\Delta \otimes \text{id})(\Delta(a)) \star (-\tilde{\varphi}), \quad a \in A, \\ \tilde{\varphi}^{1,2,3,4} \star \tilde{\varphi}^{12,3,4} &= \tilde{\varphi}^{2,3,4} \star \tilde{\varphi}^{1,23,4} \star \tilde{\varphi}^{1,2,3}, \end{aligned}$$

where we set $f \star g = f + g + \frac{1}{2}P(f, g) + \dots$, the Cambell-Baker-Hausdorff (CBH) series of the Lie algebra (A, P) .

If $\tilde{f} \in \mathfrak{m}_A^{\hat{\otimes} 2}$, we define the twist of the Drinfeld algebra $(A, m_0, P, \Delta, \tilde{\varphi})$ by $\tilde{f}$ as the algebra $(A, m_0, P, \tilde{f}\Delta, \tilde{f}\tilde{\varphi})$, where

$$\begin{aligned} \tilde{f}\Delta(a) &= \tilde{f} \star \Delta(a) \star (-\tilde{f}), \quad \text{and} \\ \tilde{f}\tilde{\varphi} &= \tilde{f}^{2,3} \star \tilde{f}^{1,23} \star \tilde{\varphi} \star (-\tilde{f}^{12,3}) \star (-\tilde{f}^{1,2}); \end{aligned}$$

then $(A, m_0, P, \tilde{f}\Delta, \tilde{f}\tilde{\varphi})$ is again a Drinfeld algebra.

Remark 1.6. If Λ is any Artinian local $\mathbb{K}$ -ring with residue field $\mathbb{K}$, set $X = \text{Hom}_{\mathbb{K}}(A, \Lambda)$. Then X is the “Poisson-Lie quasi-group”, in the sense of the Introduction. Namely, Δ_0 induces a product $m_X : X \times X \rightarrow X$, and $\exp(V_{\tilde{\varphi}})$, $\exp(V_{\tilde{\varphi}^{12,3,4}})$, etc., induce automorphisms Φ_X , $\Phi_X^{12,3,4}$, etc., of X , that satisfy the quasi-group axioms (we denote by V_f the Hamiltonian derivation of $A^{\hat{\otimes} k}$ induced by $f \in A^{\hat{\otimes} k}$). Moreover, if $\tilde{f}$ is a twist of A , then $\exp(V_{\tilde{f}})$, $\exp(V_{\tilde{f}^{12,3}})$, $\exp(V_{\tilde{f}^{1(2)3,4}})$, etc., define a twistor $(F_X, F_X^{12,3}, F_X^{(12)3,4}, \dots)$ of (X, m_X, Φ_X) . Twisting A by $\tilde{f}$ corresponds to twisting (X, m_X, Φ_X) by $(F_X, F_X^{12,3}, \dots)$.

Lemma 1.7. *If $(A, m_0, P, \Delta, \tilde{\varphi})$ is a Drinfeld algebra, set $\mathfrak{g} = \mathfrak{m}_A / (\mathfrak{m}_A)^2$; then P induces a Lie bracket μ on $\mathfrak{g}$, $\Delta - \Delta^{1,2}$ induces a linear map $\delta : \mathfrak{g} \rightarrow \Lambda^2(\mathfrak{g})$, and the reduction of $\text{Alt}(\tilde{\varphi})$ is an element φ of $\Lambda^3(\mathfrak{g})$. Then $(\mathfrak{g}, \mu, \delta, \varphi)$ is a Lie quasi-bialgebra. Moreover, twisting $(A, m_0, P, \Delta, \tilde{\varphi})$ by $\tilde{f}$ corresponds to twisting $(\mathfrak{g}, \mu, \delta, \varphi)$ by*

$$f := (\text{Alt}(\tilde{f}) \bmod (\mathfrak{m}_A)^2 \otimes \mathfrak{m}_A + \mathfrak{m}_A \otimes (\mathfrak{m}_A)^2) \in \Lambda^2(\mathfrak{g}).$$

Taking the reduction modulo $\hbar$ induces a natural map

$$\mathcal{Q}_{\text{adm}} / \mathcal{T}_{\text{adm}} \rightarrow \{\text{Drinfeld algebra structures on } \hat{S}(\mathfrak{g})\} / \text{twists}.$$

To summarize, we have a diagram

$$\begin{array}{ccc} \mathcal{Q} / \mathcal{T} & \leftarrow \mathcal{Q}_{\text{adm}} / \mathcal{T}_{\text{adm}} & \rightarrow \left\{ \begin{array}{c} \text{Drinfeld algebra structures} \\ \text{on } \hat{S}(\mathfrak{g}) \end{array} \right\} / \text{twists} \\ \text{class} \downarrow & & \downarrow \text{red} \\ & & \{\text{Lie quasi-bialgebra structures on } (\mathfrak{g}, \mu)\} / \text{twists}, \end{array}$$

where *class* is the classical limit map described in [Dr2], and *red* is the map described in Lemma 1.7. It is easy to see that this diagram commutes.

When U is a Hopf QUE algebra, the corresponding Drinfeld algebra is the Hopf-Poisson structure on $\mathcal{O}_{G^*} = (U(\mathfrak{g}^*))^*$, and $\tilde{\varphi} = 0$.

Let $(\mathfrak{g}, \mu, \delta, \varphi)$ be a Lie quasi-bialgebra. A *lift* of $(\mathfrak{g}, \mu, \delta, \varphi)$ is a Drinfeld algebra, whose reduction is $(\mathfrak{g}, \mu, \delta, \varphi)$. A general problem is to construct a lift for any Lie quasi-bialgebra. We will not solve this problem, but we will give partial existence and unicity results.

Assume that $\delta = 0$. A quasi-Lie bialgebra is then the same as a triple $(\mathfrak{g}, \mu, \varphi)$ of a Lie algebra $(\mathfrak{g}, \mu)$ and $\varphi \in \wedge^3(\mathfrak{g})^{\mathfrak{g}}$.

Theorem 1.8.

1) *There exists a lift*

$$(\hat{S}(\mathfrak{g}), m_0, P_{\mathfrak{g}^*}, \Delta_0, \tilde{\varphi}) \tag{1.2}$$

of $(\mathfrak{g}, \mu, \delta = 0, \varphi)$. Here $P_{\mathfrak{g}^}$ is the Kostant-Kirillov Poisson structure on $\mathfrak{g}^*$ and Δ_0 is the coproduct for which the elements of $\mathfrak{g}$ are primitive.*

2) *Any two lifts of $(\mathfrak{g}, \mu, \delta = 0, \varphi)$ of the form (1.2) are related by a $\mathfrak{g}$ -invariant twist.*

Examples of quasi-Lie bialgebras with $\delta = 0$ arise from *metrized Lie algebras*, i.e., pairs $(\mathfrak{g}, t_{\mathfrak{g}})$ of a Lie algebra $\mathfrak{g}$ and $t_{\mathfrak{g}} \in S^2(\mathfrak{g})^{\mathfrak{g}}$. Then $\varphi = [t_{\mathfrak{g}}^{1,2}, t_{\mathfrak{g}}^{2,3}]$. Recall that a *Lie associator* is a noncommutative formal series $\Phi(A, B)$, such that $\log \Phi(A, B)$ is a Lie series $[A, B] + \text{higher degrees terms}$, satisfying the pentagon and hexagon identities (see [Dr3]).

Proposition 1.9. *If Φ is a Lie associator, we may set $\varphi = \log(\Phi)(\bar{t}_{\mathfrak{g}}^{1,2}, \bar{t}_{\mathfrak{g}}^{2,3})$, where $\bar{t}_{\mathfrak{g}}^{i,j}$ is the image of $t_{\mathfrak{g}}^{i,j}$ in $\hat{S}(\mathfrak{g})^{\otimes 3}$, and we use the Poisson bracket of $\hat{S}(\mathfrak{g})^{\otimes 3}$ in the expression of $\log(\Phi)(\bar{t}_{\mathfrak{g}}^{1,2}, \bar{t}_{\mathfrak{g}}^{2,3})$.*

We prove these results in Section 6. If now Φ is a general (non-Lie) associator, $(U(\mathfrak{g})[[\hbar]], m_0, \Delta_0, \Phi(\hbar t_{\mathfrak{g}}^{1,2}, \hbar t_{\mathfrak{g}}^{2,3}))$ is a quasi-Hopf QUE algebra, but it is admissible only when Φ is Lie (for general $\mathfrak{g}$). According to Theorem 1.3 2), it is twist-equivalent to an admissible quasi-Hopf QUE algebra. We prove

Theorem 1.10. Any (non-Lie) associator is twist-equivalent to a unique Lie associator.

So the “concrete” version of the twist of Theorem 1.10 is an example of the twist F of Theorem 1.3 2).

§ 2 Definition and properties of U'

In this section, we prove Theorem 1.1. We first introduce the material for the definition of U' : trees (a); the map $\delta^{(P)}$ (b); then we prove Theorem 1.1 in (c) and (d).

- a - Binary complete planar rooted trees

Definition 2.1. A n -binary complete planar rooted tree (n -tree for short) is a set of vertices and oriented edges satisfying the following conditions:

- each edge carries one of the labels $\{l, r\}$.
- if we set:

$$\text{valency of a vertex} = (\text{card}(\text{incoming edges}), \text{card}(\text{outgoing edges})),$$

we have

- there exists exactly one vertex with valency $(0, 2)$ (the root)
- there exists exactly n vertices with valency $(1, 0)$ (the leaves)
- all other vertices have valency $(1, 2)$
- if a vertex has valency $(x, 2)$, then one of its outgoing edges has label l and the other has label r .
- the set of leaves has cardinal n .

Let us denote, for $n \geq 2$,

$$\text{Tree}_n = \{n\text{-binary complete planar rooted trees}\}.$$

By definition, Tree_1 consists of one element (the tree with a root and one nonmarked edge) and Tree_0 consists of one element (the tree with a root and no edge). We will write $|P| = n$ if P is a tree in Tree_n .

Definition 2.2. (Extracted trees) Let P be a binary complete planar rooted tree. Let L be the set of its leaves and let L' be a subset of L . We define the extracted subtree $P_{L'}$ as follows:

- (1) $\tilde{P}_{L'}$ is the set of all edges connecting the root with an element of L' ,
- (2) the vertices of $\tilde{P}_{L'}$ all have valency $(0, 2)$, $(1, 0)$, $(1, 2)$ or $(1, 1)$;
- (3) $P_{L'}$ is obtained from $\tilde{P}_{L'}$ by replacing each maximal sequence of edges related by a $(1, 1)$ vertex, by a single edge whose label is the label of the first edge of the sequence.

Then $P_{L'}$ is a $|L'|$ -binary complete planar rooted tree.

Definition 2.3. (Descendants of a tree) If we cut the tree P by removing its root and the related vertices, we get two trees P' and P'' , its left and right descendants.

In the same way, we define the left and right descendants of a vertex of P .

If P is a n -tree, there exists a unique bijection of the set of leaves with $\{1, \dots, n\}$, such that for each vertex, the number attached to any leaf of its left descendant is smaller than the number attached to any leaf of its right descendant.

- b - Definition of $\delta^{(P)} : U \rightarrow U^{\widehat{\otimes} n}$

Let us place ourselves in the hypothesis of Theorem 1.1. Let us define $\delta^{(2)} : U \rightarrow U^{\widehat{\otimes} 2}$,

$$\delta^{(2)}(x) = \Delta(x) - x \otimes 1 - 1 \otimes x + \varepsilon(x)1 \otimes 1.$$

For P_2 the only tree of Tree_2 , we set

$$\delta^{(P_2)} = \delta^{(2)} = \delta.$$

For P_1 , the only tree of Tree_1 , we set

$$\Delta^{(P_1)}(x) = \delta^{(1)}(x) = x - \varepsilon(x)1.$$

For P_0 the only tree of Tree_0 , we set

$$\delta^{(P_0)}(x) = \delta^{(0)}(x) = \varepsilon(x).$$

When P is a n -tree with descendants P' and P'' , we set

$$\delta^{(P)} = (\delta^{(P')} \otimes \delta^{(P'')}) \circ \delta,$$

so $\delta^{(P)}$ is a linear map $U \rightarrow U^{\widehat{\otimes} n}$.

- c - Behavior of $\delta^{(P)}$ with respect to multiplication

If $\Sigma = \{i_1, \dots, i_k\}$ is a subset of $\{1, \dots, n\}$, where $i_1 < i_2 < \dots < i_k$, the map $x \mapsto x^\Sigma$ is the linear map $U^{\widehat{\otimes} k} \rightarrow U^{\widehat{\otimes} n}$, defined by

$$x_1 \otimes \dots \otimes x_k \mapsto 1^{\otimes i_1-1} \otimes x_1 \otimes 1^{\otimes i_2-i_1-1} \otimes x_2 \otimes \dots \otimes 1^{\otimes i_k-i_{k-1}-1} \otimes x_k \otimes 1^{\otimes n-i_k-1}.$$

If $\Sigma = \emptyset$, $x \mapsto x^\Sigma$ is the map $\mathbb{K} \rightarrow U^{\widehat{\otimes} n}$, $1 \mapsto 1^{\otimes n}$.

Proposition 2.4. *For $P \in \text{Tree}_n$, we have the identity*

$$\delta^{(P)}(xy) = \sum_{\substack{\Sigma', \Sigma'' \subset \{1, \dots, n\} \\ \Sigma' \cup \Sigma'' = \{1, \dots, n\}}} (\delta^{(\Sigma')}(x))^{\Sigma'} (\delta^{(\Sigma'')}(y))^{\Sigma''},$$

for any $x, y \in U$.

This proposition is proved in Section 5.

- d - Construction of U'

Let us set

$$U' = \{x \in U \mid \text{for any tree } P, \delta^{(P)}(x) \in \hbar^{|P|} U^{\widehat{\otimes} |P|}\}.$$

Then U' is a topologically free $\mathbb{K}[[\hbar]]$ -submodule of U . Moreover, if $x, y \in U'$, and P is a tree, then

$$\delta^{(P)}([x, y]) = \sum_{\substack{\Sigma, \Sigma' \subset \{1, \dots, |P|\} \\ \Sigma \cup \Sigma' = \{1, \dots, |P|\}}} \left[\delta^{(P_\Sigma)}(x)^\Sigma, \delta^{(P_{\Sigma'})}(y)^{\Sigma'} \right];$$

the summand corresponding to a pair (Σ, Σ') with $\Sigma \cap \Sigma' = \emptyset$ is zero, and the $\hbar$ -adic valuation of the other summands is $\leq |\Sigma| + |\Sigma'| \leq |P| + 1$; so $\delta^{(P)}([x, y]) \in \hbar^{|P|+1} U^{\widehat{\otimes} |P|}$. On the other hand, there exists $z \in U$ such that $[x, y] = \hbar z$, so $\delta^{(P)}(z) \in \hbar^{|P|} U^{\widehat{\otimes} |P|}$; so $z \in U'$ and we get $[x, y] \in \hbar U'$. It follows that $U'/\hbar U'$ is commutative. Let us set

$$U'^{(n)} = U' \cap \hbar^n U. \quad (2.3)$$

We have a decreasing filtration

$$U' = U'^{(0)} \supset U'^{(1)} \supset U'^{(2)} \supset \dots;$$

we have $U'^{(n)} \subset \hbar^n U$, so U' is complete for the topology induced by this filtration. This is an algebra filtration, i.e., $U'^{(i)} U'^{(j)} \subset U'^{(i+j)}$. It induces an algebra filtration on $U'/\hbar U'$,

$$U'/\hbar U' \supset \dots \supset U'^{(i)}/(U'^{(i)} \cap \hbar U') \supset \dots,$$

for which $U'/\hbar U'$ is complete. Moreover, the completed tensor product

$$U' \bar{\otimes} U' = \varprojlim_n (U' \widehat{\otimes} U' / \sum_{p, q | p+q=n} U'^{(p)} \widehat{\otimes} U'^{(q)})$$

identifies with

$$\begin{aligned} & \varprojlim_n (\{x \in U \widehat{\otimes} U | \forall P, Q, (\delta^{(P)} \otimes \delta^{(Q)})(x) \in \hbar^{|P|+|Q|} U^{\widehat{\otimes} 2}\} / \\ & \{x \in U \widehat{\otimes} U | \forall P, Q, (\delta^{(P)} \otimes \delta^{(Q)})(x) \in \hbar^{\max(n, |P|+|Q|)} U^{\widehat{\otimes} 2}\}). \end{aligned}$$

If $x \in U'$, and P, Q are trees, with $|P|, |Q| \neq 0$, then since $\delta^{(P)}(1) = \delta^{(Q)}(1) = 0$, we have

$$\begin{aligned} (\delta^{(P)} \otimes \delta^{(Q)})(\Delta(x)) &= (\delta^{(P)} \otimes \delta^{(Q)})(\delta(x)) = \delta^{(R)}(x) \in \hbar^{|R|} U^{\widehat{\otimes} |R|} \\ &= \hbar^{|P|+|Q|} U^{\widehat{\otimes} |P|+|Q|}, \end{aligned}$$

where R is the tree whose left and right descendants are P and Q ; so $|R| = |P| + |Q|$. On the other hand,

$$\begin{aligned} (\delta^{(P)} \otimes \varepsilon)(\Delta(x)) &= \delta^{(P)}(x) \otimes 1 \in \hbar^{|P|} U^{\otimes |P|} \\ (\varepsilon \otimes \delta^{(P)})(\Delta(x)) &= 1 \otimes \delta^{(P)}(x) \in \hbar^{|P|} U^{\otimes |P|}, \end{aligned}$$

so $\Delta(x)$ satisfies $(\delta^{(P)} \otimes \delta^{(Q)})(\Delta(x)) \in \hbar^{|P|+|Q|} U^{\bar{\otimes} |P|+|Q|}$ for any pair of trees (P, Q) . $\Delta : U \rightarrow U \widehat{\otimes} U$ therefore induces an algebra morphism $\Delta_{U'} : U' \rightarrow U'^{\bar{\otimes} 2}$, whose reduction modulo $\hbar$ is a morphism of complete local rings

$$\mathcal{O} \rightarrow \mathcal{O}^{\bar{\otimes} 2} = \varprojlim_n \left(\mathcal{O}^{\otimes 2} / \sum_{p, q | p+q=n} \mathcal{O}_p \otimes \mathcal{O}_q \right),$$

where $\mathcal{O} = U' / \hbar U'$ and $\mathcal{O}_p = U'^{(p)} / (U'^{(p)} \cap \hbar U')$.

§ 3 Classical limit of U'

We will prove Theorem 1.2 as follows. We first compare the various $\delta^{(P)}$, where P is a n -tree (Proposition 3.1). Relations found between the $\delta^{(P)}$ imply that they have $\hbar$ -adic valuation properties close to those of the Hopf case (Proposition 3.2). We then prove Theorem 1.2.

- a - Comparison of the various $\delta^{(P)}$

Let P and P_0 be n -trees. There exists an element $\Phi^{P,P_0} \in U^{\otimes n}$, such that $\Delta^{(P)} = \text{Ad}(\Phi^{P,P_0}) \circ \Delta^{(P_0)}$. The element Φ^{P,P_0} is a product of images of Φ and Φ^{-1} by the various maps $U^{\otimes 3} \rightarrow U^{\otimes n}$ obtained by iteration of Δ . We have

$$\Phi^{P',P_0} = \Phi^{P',P} \Phi^{P,P_0} \quad (3.4)$$

for any n -trees P_0, P, P' . For example,

$$\begin{aligned} (\text{id} \otimes \Delta) \circ \Delta &= \text{Ad}(\Phi) \circ ((\Delta \otimes \text{id}) \circ \Delta), \\ (\Delta \otimes \Delta) \circ \Delta &= \text{Ad}(\Phi^{12,3,4}) \circ ((\Delta \otimes \text{id}^{\otimes 2}) \circ (\Delta \otimes \text{id}) \circ \Delta), \text{ etc.} \end{aligned}$$

Proposition 3.1. Assume that $\hbar \log(\Phi) \in (U')^{\bar{\otimes} 3}$. Then there exists a sequence of elements

$$F^{PP_0R\Sigma v} = \sum_{\alpha} F_{1,\alpha}^{PP_0R\Sigma v} \otimes \cdots \otimes F_{v,\alpha}^{PP_0R\Sigma v} \in (U'^{\bar{\otimes} n})^{\bar{\otimes} v},$$

indexed by the triples (R, Σ, v) , where R is a tree such that $|R| < n$, Σ is a subset of $\{1, \dots, n\}$ with $\text{card}(\Sigma) = |R|$, and v is an integer ≥ 1 , such that the equality

$$\begin{aligned} \delta^{(P)} &= \text{Ad}(\Phi^{P,P_0}) \circ \delta^{(P_0)} + \sum_{k|k < n} \sum_{R \text{ a } k\text{-tree}} \sum_{\substack{\Sigma \subset \{1, \dots, n\}, \\ \text{card}(\Sigma) = k}} \\ &\quad \sum_{v \geq 1} \sum_{\alpha} \text{ad}_{\hbar}(F_{1,\alpha}^{PP_0R\Sigma v}) \circ \cdots \circ \text{ad}_{\hbar}(F_{v,\alpha}^{PP_0R\Sigma v}) \circ (\delta^{(R)})^{\Sigma} \end{aligned} \quad (3.5)$$

holds. Here $\text{ad}_{\hbar}(x)(y) = \frac{1}{\hbar} [x, y]$.

PROOF. Let us prove this statement by induction on n . When $n = 3$, we find

$$\delta^{(1(23))} = \text{Ad}(\Phi) \delta^{((12)3)} + (\text{Ad}(\Phi) - 1)(\delta^{1,2} + \delta^{1,3} + \delta^{2,3} + \delta^{(1)1} + \delta^{(1)2} + \delta^{(1)3}),$$

so the identity holds with $F^{PP_0R\Sigma v} = \frac{1}{v!} (\hbar \log \Phi)^{\bar{\otimes} v}$ for all choices of (R, Σ, v) , except when $|R| = 0$, in which case $F^{PP_0R\Sigma v} = 0$. Assume that the statement holds for any pair of k -trees, $k \leq n$, and let us prove it for a pair (P, P_0) of $(n+1)$ -trees. For k any integer, let $P_{\text{left}}(k)$ be the k -tree corresponding to

$$\delta^{(P_{\text{left}}(k))} = (\delta \otimes \text{id}^{\otimes k-2}) \circ \cdots \circ \delta.$$

Thanks to (3.4), we may assume that $P_0 = P_{\text{left}}(n+1)$ and P is arbitrary. Let P' and P'' be the subtrees of P , such that $|P'| + |P''| = n+1$, and $\delta^{(P)} = (\delta^{(P')} \otimes \delta^{(P'')}) \circ \delta$. Let P_1 and P_2 the n -trees such that

$$\delta^{(P_1)} = (\delta^{(P_{\text{left}}(k'))} \otimes \delta^{(P'')}) \circ \delta \text{ and } \delta^{(P_2)} = (\delta^{(P_{\text{left}}(k'))} \otimes \delta^{(P_{\text{left}}(k''))}) \circ \delta$$

Assume that $|P_1| \neq 1$. Using (3.4), we reduce the proof of (3.5) to the case of the pairs (P, P_1) , (P_1, P_2) and (P_2, P_0) . Then the induction hypothesis applied to the pair $(P', P_{\text{left}}(k'))$, together with $\Phi^{P, P_1} = \Phi^{P', P_{\text{left}}(k')} \otimes 1^{\otimes k''}$, implies

$$\begin{aligned} \delta^{(P)} &= \text{Ad}(\Phi^{P, P_1}) \circ \delta^{(P_1)} + \sum_{k|k < k' \text{ a } k\text{-tree}} \sum_{\substack{\Sigma \subset \{1, \dots, k'\}, \\ \text{card}(\Sigma) = k}} \sum \\ &\quad \sum_{v \geq 1} \sum_{\alpha} \text{Ad}(\Phi^{P, P_1}) \circ \text{ad}_{\bar{h}}(F_{1, \alpha}^{P' P_{\text{left}}(k') \Sigma v} \otimes 1^{\otimes k''}) \dots \text{ad}_{\bar{h}}(F_{v, \alpha}^{P' P_{\text{left}}(k') \Sigma v} \otimes 1^{\otimes k''}) \\ &\quad \circ ((\delta^{(R)} \otimes \delta^{(P'')}) \circ \delta)^{\Sigma, k'+1, \dots, n+1}, \end{aligned}$$

which is (3.5) for (P, P_1) . In the same way, one proves a similar identity relating P_1 and P_2 . Let us now prove the identity relating P_2 and P_0 . We have $\delta^{(P_2)} = (\delta \otimes \text{id}^{\otimes n-1}) \circ \delta^{(P'_2)}$ and $\delta^{(P_0)} = (\delta \otimes \text{id}^{\otimes n-1}) \circ \delta^{(P'_0)}$, where P'_2 and P'_0 are n -trees. We have

$$\Phi^{P_2, P_0} = (\Delta \otimes \text{id}^{\otimes n-2}) \circ \Phi^{P'_2, P'_0}$$

so we get

$$\begin{aligned} \delta^{(P_2)} &= \text{Ad}(\Phi^{P_2, P_0}) \circ \delta^{(P_0)} \\ &+ (\text{Ad}(\Phi^{P_2, P_0}) - \text{Ad}((\Phi^{P'_2, P'_0})^{1, 3, \dots, n+1})) \circ (\delta^{(P'_0)})^{1, 3, \dots, n+1} \\ &+ (\text{Ad}(\Phi^{P_2, P_0}) - \text{Ad}((\Phi^{P'_2, P'_0})^{2, 3, \dots, n+1})) \circ (\delta^{(P'_0)})^{2, 3, \dots, n+1} \\ &+ (\delta \otimes \text{id}^{\otimes n-1}) \left(\sum_{k \leq n} \sum_{\text{a } k\text{-tree}} \sum_{\substack{\Sigma \subset \{1, \dots, n\}, \\ \text{card}(\Sigma) = k}} \right. \\ &\quad \left. \sum_{v \geq 1} \sum_{\alpha} \text{ad}_{\bar{h}}(F_{1, \alpha}^{P'_2 P'_0 \Sigma v}) \dots \text{ad}_{\bar{h}}(F_{v, \alpha}^{P'_2 P'_0 \Sigma v}) \circ (\delta^{(R)})^{\Sigma} \right). \end{aligned}$$

We have $\bar{h} \log \Phi^{P_2, P_0} \in U'^{\bar{\otimes} n+1}$ and $\bar{h} \log \Phi^{P'_2, P'_0} \in U'^{\bar{\otimes} n}$; this fact and the relations

$$\begin{aligned} (\delta \otimes \text{id}^{\otimes n-1}) (\text{ad}_{\bar{h}}(x_1) \dots \text{ad}_{\bar{h}}(x_v) \circ (\delta^{(R)})^{\Sigma}) &= \\ (\text{ad}_{\bar{h}}(x_1^{12, \dots, n+1}) \circ \dots \circ \text{ad}_{\bar{h}}(x_v^{12, \dots, n+1}) - \text{ad}_{\bar{h}}(x_1^{1, 3, \dots, n+1}) \circ \dots \circ \text{ad}_{\bar{h}}(x_v^{1, 3, \dots, n+1}) \\ &\quad - \text{ad}_{\bar{h}}(x_1^{2, 3, \dots, n+1}) \circ \dots \circ \text{ad}_{\bar{h}}(x_v^{2, 3, \dots, n+1})) \circ (\delta^{(R)})^{\Sigma+1} \end{aligned}$$

if $1 \notin \Sigma$, and

$$\begin{aligned} (\delta \otimes \text{id}^{\otimes n-1}) (\text{ad}_{\bar{h}}(x_1) \dots \text{ad}_{\bar{h}}(x_v) \circ (\delta^{(R)})^{\Sigma}) &= \\ \text{ad}_{\bar{h}}(x_1^{12, \dots, n+1}) \circ \dots \circ \text{ad}_{\bar{h}}(x_v^{12, \dots, n+1}) \circ ((\delta \otimes \text{id}^{\otimes n-1}) \circ \delta^{(R)})^{1, 2, \Sigma'+1} \\ &+ (\text{ad}_{\bar{h}}(x_1^{12, \dots, n+1}) \circ \dots \circ \text{ad}_{\bar{h}}(x_v^{12, \dots, n+1}) - \text{ad}_{\bar{h}}(x_1^{1, 3, \dots, n+1}) \circ \dots \circ \text{ad}_{\bar{h}}(x_v^{1, 3, \dots, n+1})) \\ &\quad \circ (\delta^{(R)})^{1, \Sigma'+1} \\ &+ (\text{ad}_{\bar{h}}(x_1^{12, \dots, n+1}) \circ \dots \circ \text{ad}_{\bar{h}}(x_v^{12, \dots, n+1}) - \text{ad}_{\bar{h}}(x_1^{2, 3, \dots, n+1}) \circ \dots \circ \text{ad}_{\bar{h}}(x_v^{2, 3, \dots, n+1})) \\ &\quad \circ (\delta^{(R)})^{2, \Sigma'+1}. \end{aligned}$$

if $\Sigma = \Sigma' \cup \{1\}$, where $1 \notin \Sigma'$, imply that $\delta^{(P_2)} - \text{Ad}(\Phi^{P_2, P_0}) \circ \delta^{(P_0)}$ has the desired form. Let us now treat the case $|P_1| = 1$. For this, we introduce the trees P_3 and P_4 , such that:

$$\begin{aligned}\delta^{(P_3)} &= (\text{id}^{\otimes n-1} \otimes \delta) \circ (\text{id}^{\otimes n-2} \otimes \delta) \circ \dots \circ \delta, \\ \delta^{(P_4)} &= (\text{id}^{\otimes n-1} \otimes \delta) \circ (\delta \otimes \text{id}^{\otimes n-2}) \circ (\delta \otimes \text{id}^{\otimes n-3}) \circ \dots \circ (\delta \otimes \text{id}) \circ \delta.\end{aligned}$$

We then prove the relation for the pair (P, P_3) in the same way as for (P_1, P_2) (only the right branch of the tree is changed); the relation for (P_3, P_4) in the same way as for (P_2, P_3) (instead of composing a known relation by $\delta \otimes \text{id}^{\otimes n-1}$, we compose it with $\text{id}^{\otimes n-1} \otimes \delta$); and using the identity

$$\delta^{(P_4)} = (\delta \otimes \text{id}^{\otimes n-1}) \circ (\text{id}^{\otimes n-2} \otimes \delta) \circ (\delta \otimes \text{id}^{\otimes n-3}) \circ \dots \circ \delta,$$

we prove the relation for (P_4, P) in the same way as for (P_2, P_3) (composing a known relation by $\delta \otimes \text{id}^{\otimes n-1}$). $\square$

- b - Properties of $\delta^{(P)}$

Proposition 3.2. *Let n be an integer and $x \in U$.*

- 1) *Assume that for any tree R , such that $|R| < n$, we have $\delta^{(R)}(x) \in \hbar^{|R|} U^{\widehat{\otimes} |R|}$. Then the conditions*

$$\delta^{(P)}(x) \in \hbar^n U^{\widehat{\otimes} n} \quad (3.6)$$

where P is an n -tree, are all equivalent.

- 2) *Assume that for any tree R , such that $|R| < n$, we have $\delta^{(R)}(x) \in \hbar^{|R|+1} U^{\widehat{\otimes} |R|}$. Then the elements*

$$\left(\frac{1}{\hbar^n} \delta^{(P)}(x) \bmod \hbar \right) \in U(\mathfrak{g})^{\otimes n},$$

where P is an n -tree, are all equal and belong to $(\mathfrak{g}^{\otimes n})^{\mathfrak{S}_n} = S^n(\mathfrak{g})$.

PROOF. Let us prove 1). We have $\delta^{(P)} = (\text{id} - \eta \circ \varepsilon)^{\otimes |P|} \circ \delta^{(P)}$, where $\eta : \mathbb{K}[[\hbar]] \rightarrow U$ is the unit map of U , so

$$\begin{aligned}\delta^{(P)} &= \text{Ad}(\Phi^{P, P_0}) \circ \delta^{(P_0)} + \sum_{k|k < n} \sum_{R \text{ a } k\text{-tree}} \sum_{\substack{\Sigma \subset \{1, \dots, n\}, \\ \text{card}(\Sigma) = k}} \sum_{v \geq 1} \sum_{\alpha} \\ &(\text{id} - \eta \circ \varepsilon)^{\otimes n} \circ \text{ad}_{\hbar}(F_{1, \alpha}^{P P_0 R \Sigma v}) \circ \dots \circ \text{ad}_{\hbar}(F_{v, \alpha}^{P P_0 R \Sigma v}) \circ (\delta^{(R)})^{\Sigma}.\end{aligned}$$

Then 1) follows from:

Lemma 3.3. *Let Σ be a subset of $\{1, \dots, n\}$ (we will write $|\Sigma|$ instead of $\text{card}(\Sigma)$) and let U_0 be the kernel of the counit of U . Let $x \in \hbar^{|\Sigma|} (U_0)^{\widehat{\otimes} |\Sigma|}$ and $F_1, \dots, F_v$ be elements of $(U')^{\otimes n}$. Then*

$$(\text{id} - \eta \circ \varepsilon)^{\otimes n} (\text{ad}_{\hbar}(F_1) \dots \text{ad}_{\hbar}(F_v)(x^{\Sigma})) \in \hbar^n (U_0)^{\widehat{\otimes} n}.$$

PROOF OF LEMMA. Each element $F \in (U')^{\widehat{\otimes} n}$ is uniquely expressed as a sum $F = \sum_{\Sigma \in \mathcal{P}(\{1, \dots, n\})} F_{\Sigma}$, where F_{Σ} belongs to the image of

$$\begin{aligned}(U_0')^{\widehat{\otimes} |\Sigma|} &\rightarrow (U')^{\widehat{\otimes} n}, \\ f &\mapsto f^{\Sigma},\end{aligned}$$

$\mathcal{P}(\{1, \dots, n\})$ is the set of subsets of $\{1, \dots, n\}$, and U'_0 is the kernel of the counit of U' . Then

$$\begin{aligned} & (\text{id} - \eta \circ \varepsilon)^{\otimes n} (\text{ad}_{\bar{h}}(F_1) \cdots \text{ad}_{\bar{h}}(F_v)(x^\Sigma)) \\ &= \sum_{\Sigma_1, \dots, \Sigma_v \in \mathcal{P}(\{1, \dots, n\})} (\text{id} - \eta \circ \varepsilon)^{\otimes n} (\text{ad}_{\bar{h}}((F_1)_{\Sigma_1}) \cdots \text{ad}_{\bar{h}}((F_v)_{\Sigma_v})(x^\Sigma)). \end{aligned}$$

The summands corresponding to $(\Sigma_1, \dots, \Sigma_v)$ such that $\Sigma_1 \cup \dots \cup \Sigma_v \cup \Sigma \neq \{1, \dots, n\}$ are all zero. Moreover, each $(F_\alpha)_{\Sigma_\alpha}$ can be expressed as $(f_\alpha)^{\Sigma_\alpha}$, where $f_\alpha \in \bar{h}^{|\Sigma_\alpha|}(U_0)^{\hat{\otimes} |\Sigma_\alpha|}$. The lemma then follows from the statement:

Statement 3.4. *If $\Sigma, \Sigma' \subset \{1, \dots, n\}$, $x \in \bar{h}^{|\Sigma|}(U_0)^{\hat{\otimes} |\Sigma|}$, $y \in \bar{h}^{|\Sigma'|}(U_0)^{\hat{\otimes} |\Sigma'|}$, then $\frac{1}{\bar{h}}[x, y]$ can be expressed as $z^{\Sigma \cup \Sigma'}$, where $z \in \bar{h}^{|\Sigma \cup \Sigma'|}(U_0)^{\hat{\otimes} |\Sigma \cup \Sigma'|}$.*

PROOF. If $\Sigma \cap \Sigma' = \emptyset$, then $[x, y] = 0$, so the statement holds. If $\Sigma \cap \Sigma' \neq \emptyset$, then the $\bar{h}$ -adic valuation of $\frac{1}{\bar{h}}[x, y]$ is $\geq -1 + |\Sigma| + |\Sigma'| \geq |\Sigma| + |\Sigma'| - |\Sigma \cap \Sigma'| = |\Sigma \cup \Sigma'|$. $\square$

Let us now prove property 2). The above arguments immediately imply that the $(\frac{1}{\bar{h}}\delta^{(P)}(x) \bmod \bar{h})$, $|P| = n$, are all equal. This defines an element $S_n(x) \in U(\mathfrak{g})^{\otimes n}$. If $|P| = n$, we have $(\text{id}^{\otimes k} \otimes \delta \otimes \text{id}^{\otimes n-k-1}) \circ \delta^{(P)}(x) \in \bar{h}^{n+1}U^{\hat{\otimes} n+1}$, so if $\delta_0 : U(\mathfrak{g}) \rightarrow U(\mathfrak{g}) \otimes U(\mathfrak{g})$ is defined by $\delta_0(x) = \Delta_0(x) - x \otimes 1 - 1 \otimes x + \varepsilon(x)1 \otimes 1$, Δ_0 being the coproduct of $U(\mathfrak{g})$, then $(\text{id}^{\otimes k} \otimes \delta_0 \otimes \text{id}^{\otimes n-k-1})(S_n(x)) = 0$, so

$$S_n \in \mathfrak{g}^{\otimes n}. \quad (3.7)$$

Let us denote by $\sigma_{i,i+1}$ the permutation of the factors i and $i+1$ in a tensor power. For $i = 1, \dots, n-1$, let us compute $(\sigma_{i,i+1} - \text{id})(S_n(x))$. Let P' be a $(n-1)$ -tree and let P be the n -tree such that $\delta^{(P)} = (\text{id}^{\otimes i-1} \otimes \delta \otimes \text{id}^{\otimes n-i-1}) \circ \delta^{(P')}$. Then

$$(\sigma_{i,i+1} - \text{id})(S_n) = \left[\frac{1}{\bar{h}} (\text{id}^{\otimes i-1} \otimes (\delta^{2,1} - \delta) \otimes \text{id}^{\otimes n-i-1}) \circ \delta^{(P')}(x) \bmod \bar{h} \right].$$

By assumption, $\delta^{(P')}(x) \in \bar{h}^n U^{\hat{\otimes} n-1}$; moreover, $\delta^{2,1} - \delta = \Delta^{2,1} - \Delta$, so $(\delta^{2,1} - \delta)(U) \subset \bar{h}(U \hat{\otimes} U)$; therefore

$$(\text{id}^{\otimes i-1} \otimes (\delta^{2,1} - \delta) \otimes \text{id}^{\otimes n-i-1}) \circ \delta^{(P')}(x) \in \bar{h}^{n+1}U^{\hat{\otimes} n},$$

it follows that $(\sigma_{i,i+1} - \text{id})(S_n(x)) = 0$, therefore $S_n(x)$ is a symmetric tensor of $U(\mathfrak{g})^{\otimes n}$. Together with (3.7), this gives $S_n(x) \in (\mathfrak{g}^{\otimes n})^{\mathfrak{S}_n}$. This ends the proof of Proposition 3.2. $\square$

- c - Flatness of U' (proof of Theorem 1.2)

Let us set

$$U''^{(n)} = \{x \in U' \mid \delta^{(P)}(x) \in \bar{h}^{|P|+1}U^{\hat{\otimes} |P|} \text{ if } |P| \leq n\}.$$

Then by Proposition 2.4, we have a decreasing algebra filtration

$$U' = U''^{(0)} \supset U''^{(1)} \supset U''^{(2)} \supset \dots \supset \bar{h}U'. \quad (3.8)$$

Each $U''^{(n)}$ is divisible in U' , i.e., $U''^{(n)} \cap \bar{h}U' = \bar{h}U''^{(n)}$. We also have $U''^{(n)} \supset U'^{(n)} + \bar{h}U'$ (we will see later that this is an equality). We derive from (3.8) a decreasing filtration

$$\mathcal{O} = \mathcal{O}''^{(0)} \supset \mathcal{O}''^{(1)} \supset \mathcal{O}''^{(2)} \supset \dots,$$

where $\mathcal{O} = U'/\hbar U'$ and $\mathcal{O}''^{(n)} = U''^{(n)}/\hbar U''^{(n)}$. We have clearly

$$\bigcap_{n \geq 0} \mathcal{O}''^{(n)} = \{0\};$$

the fact that $\mathcal{O}$ is complete for this filtration will follow from its identification with the filtration $\mathcal{O} \supset \mathcal{O}'^{(1)} \supset \dots$ (see Proposition 3.6), where $\mathcal{O}'^{(i)} = U'^{(i)}/\hbar U \cap U'^{(i)}$ and $U'^{(i)}$ is defined in (2.3). We first prove:

Proposition 3.5. *Set $\widehat{\text{gr}}''(\mathcal{O}) = \bigoplus_{n \geq 0} \mathcal{O}''^{(n)}/\mathcal{O}''^{(n+1)}$. Then there is a unique linear map $\lambda_n : \text{gr}_n''(\mathcal{O}) \rightarrow S^n(\mathfrak{g})$, taking the class of x to the common value of all $\frac{1}{n!}(\frac{1}{\hbar^n} \delta^{(P)}(x) \bmod \hbar)$, where P is a n -tree. The resulting map $\lambda : \widehat{\text{gr}}''(\mathcal{O}) \rightarrow \widehat{S}(\mathfrak{g})$ is an isomorphism of graded complete algebras.*

PROOF. In Proposition 3.2, we constructed a map $U''^{(n)} \rightarrow S^n(\mathfrak{g})$, by $x \mapsto$ common value of $\frac{1}{n!}(\frac{1}{\hbar^n} \delta^{(P)}(x) \bmod \hbar)$ for all n -trees P . The subspace $U''^{(n+1)} \subset U''^{(n)}$ is clearly contained in the kernel of this map, so we obtain a map

$$\lambda_n : U''^{(n)}/U''^{(n+1)} = \mathcal{O}''^{(n)}/\mathcal{O}''^{(n+1)} \rightarrow S^n(\mathfrak{g}).$$

Let us prove that $\lambda = \bigoplus_{n \geq 1} \lambda_n$ is a morphism of algebras. If $x \in U''^{(n)}$ and $y \in U''^{(m)}$, Proposition 2.4 implies that if R is any $(n+m)$ -tree, we have

$$\delta^{(P)}(xy) = \sum_{\substack{\Sigma', \Sigma'' \subset \{1, \dots, n+m\} \\ \Sigma' \cup \Sigma'' = \{1, \dots, n+m\}}} \delta^{(R_{\Sigma'})}(x)^{\Sigma'} \delta^{(R_{\Sigma''})}(y)^{\Sigma''}.$$

The $\hbar$ -adic valuation of the term corresponding to (Σ', Σ'') is $\geq |\Sigma'| + |\Sigma''|$ if $|\Sigma'| \geq n$ and $|\Sigma''| \geq m$, and $\geq |\Sigma'| + |\Sigma''| + 1$ otherwise, so the only contributions to $(\frac{1}{\hbar^{n+m}} \delta^{(R)}(xy) \bmod \hbar)$ are those of the pairs (Σ', Σ'') such that $\Sigma' \cap \Sigma'' = \emptyset$. Then:

$$\begin{aligned} & \left(\frac{1}{\hbar^{n+m}} \delta^{(R)}(xy) \bmod \hbar \right) \\ &= \sum_{\substack{\Sigma', \Sigma'' \subset \{1, \dots, n+m\} \\ |\Sigma'|=n, |\Sigma''|=m, \\ \Sigma' \cap \Sigma'' = \emptyset}} \left(\frac{1}{\hbar^n} \delta^{(R_{\Sigma'})}(x) \bmod \hbar \right) \left(\frac{1}{\hbar^m} \delta^{(R_{\Sigma''})}(y) \bmod \hbar \right) \\ &= \sum_{\substack{\Sigma', \Sigma'' \subset \{1, \dots, n+m\} \\ |\Sigma'|=n, |\Sigma''|=m, \\ \Sigma' \cap \Sigma'' = \emptyset}} (n! \lambda_n(x)^{\Sigma'}) (m! \lambda_m(y)^{\Sigma''}) \\ &= (n+m)! \lambda_n(x) \lambda_m(y), \end{aligned}$$

because the map

$$\begin{aligned} S(\mathfrak{g}) &\rightarrow (T(\mathfrak{g}), \text{shuffle product}), \\ x_1 \cdots x_n &\mapsto \sum_{\sigma \in \mathfrak{S}_n} x_{\sigma(1)} \otimes \cdots \otimes x_{\sigma(n)} \end{aligned}$$

is an algebra morphism. Therefore $\lambda_{n+m}(xy) = \lambda_n(x)\lambda_m(y)$. Let us prove that λ_n is injective. If $x \in U''^{(n)}$ is such that $(\frac{1}{h^n}\delta^{(P)}(x) \bmod \bar{h}) = 0$ for any n -tree P , then $x \in U''^{(n+1)}$, so its class in $\mathcal{O}''^{(n)}/\mathcal{O}''^{(n+1)} = U''^{(n)}/U''^{(n+1)}$ is zero. So each λ_n is injective, so λ is injective.

To prove that λ is surjective, it suffices to prove that λ_1 is surjective. Let us fix $x \in \mathfrak{g}$. We will construct a sequence $x_n \in U$, $n \geq 0$ such that $\varepsilon(x_n) = 0$, $(\frac{1}{h}x_n \bmod \bar{h}) = x$, $x_{n+1} \in x_n + \bar{h}^{n+1}U$ for any $n \geq 1$, and if P is any tree such that $|P| \leq n$, $\delta^{(P)}(x_n) \in \bar{h}^{|P|}U^{\widehat{\otimes}|P|}$ (this last condition implies that $\delta^{(Q)}(x_n) \in \bar{h}^n U^{\widehat{\otimes}|Q|}$ for $|Q| \geq n$). Then the limit $\tilde{x} = \lim_{n \rightarrow \infty} (x_n)$ exists, belongs to U' , satisfies $\varepsilon(\tilde{x}) = 0$ and $(\frac{1}{h}\delta_1(\tilde{x}) \bmod \bar{h}) = x$, so its class in $U''^{(1)}/U''^{(2)}$ is a preimage of x .

Let us now construct the sequence $(x_n)_{n \geq 0}$. We fix a linear map $\mathfrak{g} \rightarrow \{y \in U \mid \varepsilon(y) = 0\}$, $y \mapsto \bar{y}$, such that for any $y \in \mathfrak{g}$, $(\bar{y} \bmod \bar{h}) = y$. We set $x_1 = \bar{h}\bar{x}$. Let us construct x_{n+1} knowing x_n . By Proposition 3.2, if Q is any $(n+1)$ -tree, $\delta^{(Q)}(x_n) \in \bar{h}^n U^{\widehat{\otimes}n+1}$, and $(\frac{1}{h^n}\delta^{(Q)}(x_n) \bmod \bar{h})$ is an element of $S^{n+1}(\mathfrak{g})$, independent of Q . Let us write this element as

$$\sum_{\sigma \in \mathfrak{S}_{n+1}} \sum_{\alpha} y_{\sigma(1)}^{\alpha} \cdots y_{\sigma(n+1)}^{\alpha}, \text{ where } \sum_{\alpha} y_1^{\alpha} \otimes \cdots \otimes y_{n+1}^{\alpha} \in \mathfrak{g}^{\otimes n+1}.$$

Then we set

$$x_{n+1} = x_n - \frac{\bar{h}^{n+1}}{(n+1)!} \sum_{\sigma \in \mathfrak{S}_{n+1}} \bar{y}_{\sigma(1)}^{\alpha} \cdots \bar{y}_{\sigma(n+1)}^{\alpha}. \quad \square$$

We now prove:

Proposition 3.6.

- 1) For any $n \geq 0$, $U''^{(n)} = U'^{(n)} + \bar{h}U'$;
- 2) The filtrations $\mathcal{O} = \mathcal{O}'^{(0)} \supset \mathcal{O}'^{(1)} \supset \cdots$ and $\mathcal{O} = \mathcal{O}''^{(0)} \supset \mathcal{O}''^{(1)} \supset \cdots$ coincide, and $\mathcal{O}$ is complete and separated for this filtration.

PROOF. Let us prove 1). We have to show that $U''^{(n)} \subset U'^{(n)} + \bar{h}U'$. Let $x \in U''^{(n)}$. We have $\delta^{(P)}(x) \in \bar{h}^{|P|+1}U^{\widehat{\otimes}|P|}$ for $|P| \leq n-1$, and for P an n -tree, $(\frac{1}{h^n}\delta^{(P)}(x) \bmod \bar{h}) \in S^n(\mathfrak{g})$ and is independent on P . Write this element of $S^n(\mathfrak{g})$ as $\sum_{\sigma \in \mathfrak{S}_n} \sum_{\alpha} y_{\sigma(1)}^{\alpha} \otimes \cdots \otimes y_{\sigma(n)}^{\alpha}$ and set $f_n = \frac{1}{n!} \sum_{\sigma \in \mathfrak{S}_n} \sum_{\alpha} \bar{y}_{\sigma(1)}^{\alpha} \cdots \bar{y}_{\sigma(n)}^{\alpha}$. Then each $\bar{y}_i^{\alpha}$ belongs to $U' \cap \bar{h}U$, so $f_n \in U' \cap \bar{h}^n U = U'^{(n)}$. Moreover, $x - f_n$ belongs to $U''^{(n+1)}$. Iterating this procedure, we construct elements $f_{n+1}, f_{n+2}, \dots$, where each f_k belongs to $U'^{(k)}$. The series $\sum_{k \geq n} f_k$ converges in U' ; denote by f its sum, then $x - f$ belongs to $\bigcap_{k \geq n} U''^{(k)} = \bar{h}U'$. So $U''^{(n)} \subset U'^{(n)} + \bar{h}U'$. The inverse inclusion is obvious. This proves 1). Then 1) immediately implies that for any n , $\mathcal{O}'^{(n)} = \mathcal{O}''^{(n)}$. We already know $\mathcal{O}$ is complete and separated for $\mathcal{O} = \mathcal{O}'^{(0)} \supset \mathcal{O}'^{(1)} \supset \cdots$, which proves 2). $\square$

END OF PROOF OF THEOREM 1.2. $\mathcal{O}$ is a complete local ring, and we have a ring isomorphism $\widehat{\text{gr}}(\mathcal{O}) \rightarrow \widehat{S}(\mathfrak{g})$. Then any lift $\mathfrak{g} \rightarrow \mathcal{O}'^{(1)}$ of $\mathcal{O}'^{(1)} \rightarrow \mathcal{O}'^{(1)}/\mathcal{O}'^{(2)} = \mathfrak{g}$ yields a continuous ring morphism $\mu : \widehat{S}(\mathfrak{g}) \rightarrow \mathcal{O}$. The associated graded of μ is the identity, so μ is an isomorphism. So $\mathcal{O}$ is noncanonically isomorphic to $\widehat{S}(\mathfrak{g})$. $\square$

Remark 3.7. When U is Hopf and $\mathfrak{g}$ is finite-dimensional, $U'/\hbar U'$ identifies canonically with $\mathcal{O}_{G^*} = (U(\mathfrak{g}^*))^*$, where $\mathfrak{g}^*$ is the dual Lie bialgebra of $\mathfrak{g}$ (see [Dr1], [Gal]). The natural projection $T(\mathfrak{g}^*) \rightarrow U(\mathfrak{g}^*)$ and the identification $T(\mathfrak{g}^*)^* = \widehat{T}(\mathfrak{g})$ (where $\widehat{T}(\mathfrak{g})$ means the degree completion) induce an injection $U'/\hbar U' = \mathcal{O}_{G^*} = (U(\mathfrak{g}^*))^* \hookrightarrow \widehat{T}(\mathfrak{g})$. The map $U'/\hbar U' \hookrightarrow \widehat{T}(\mathfrak{g})$ can be interpreted simply as follows. For any $x \in U'$, we have $(\frac{1}{\hbar} \delta_n(x) \bmod \hbar) \in \mathfrak{g}^{\otimes n}$. Then $U'/\hbar U' \hookrightarrow \widehat{T}(\mathfrak{g})$ takes the class of $x \in U'$ to the sequence $(\frac{1}{\hbar} \delta_n(x) \bmod \hbar)_{n \geq 0}$.

In the quasi-Hopf case, we have no canonical embedding $U'/\hbar U' \hookrightarrow \widehat{T}(\mathfrak{g})$ because the various $(\frac{1}{\hbar} \delta^{(P)}(x) \bmod \hbar)$ do not necessarily coincide for all the n -trees P . This is related to the fact that one cannot expect a Hopf pairing $U(\mathfrak{g}^*) \otimes (U'/\hbar U') \rightarrow \mathbb{K}$ since $\mathfrak{g}^*$ is no longer a Lie algebra, so $U(\mathfrak{g}^*)$ does not make sense.

In the other hand, Theorem 1.2 can be interpreted as follows: in the Hopf case, the exponential induces an isomorphism of formal schemes $\mathfrak{g}^* \rightarrow G^*$, so $U'/\hbar U'$ identifies noncanonically with $\mathcal{O}_{G^*} = \widehat{S}(\mathfrak{g})$. In the quasi-Hopf case, although there is no formal group G^* , we still have an isomorphism $U'/\hbar U' \xrightarrow{\sim} \widehat{S}(\mathfrak{g})$. $\square$

§ 4 Twists

- a - Admissible twists

If (U, m, Δ, Φ) is an arbitrary QHQUE algebra, we will call a twist $F \in (U^{\widehat{\otimes} 2})^\times$ *admissible* if $\hbar \log(F) \in (U')^{\widehat{\otimes} 2}$.

Proposition 4.1. *Let (U, m, Δ, Φ) be an admissible quasi-Hopf algebra and F an admissible twist. Then the twisted quasi-Hopf algebra $(U, m, {}^F\Delta, {}^F\Phi)$ is admissible.*

PROOF. Set $f = \hbar \log(F)$. Then we have

$$\hbar \log({}^F\Phi) = f^{1,2} \star f^{12,3} \star (\hbar \log(\Phi)) \star (-f^{1,23}) \star (-f^{2,3}),$$

where $a \star b = a + b + \frac{1}{\hbar}[a, b] + \dots$ (the CBH series for $U'^{\widehat{\otimes} 3}$ equipped with the bracket $\frac{1}{\hbar}[-, -]$). Since $U'^{\widehat{\otimes} 3}$ is stable under $\star$, we have $\hbar \log({}^F\Phi) \in U'^{\widehat{\otimes} 3}$. So $(U, m, {}^F\Delta, {}^F\Phi)$ is admissible. $\square$

Let us now prove

Proposition 4.2. *Under the hypothesis of Proposition 4.1, the QFS algebra U'_F corresponding to $(U, m, {}^F\Delta, {}^F\Phi)$ coincides with the QFS algebra U' corresponding to (U, m, Δ, Φ) .*

We will first prove the following lemma:

Lemma 4.3. *Let P be an n -tree. Then*

$$\begin{aligned} \delta_F^{(P)} &= \delta^{(P)} + \sum_{k \leq nR} \sum_{a \text{ a } k\text{-tree}} \sum_{\substack{\Sigma \subset \{1, \dots, n\} \\ \text{card}(\Sigma) = k}} \\ &\quad \sum_{v \geq 1} \sum_{\alpha} \text{ad}_{\hbar}(f_{1, \alpha}^{\Sigma, P}) \circ \dots \circ \text{ad}_{\hbar}(f_{v, \alpha}^{\Sigma, P}) \circ (\delta^{(R)})^{\Sigma}, \end{aligned} \tag{4.9}$$

where for each v , $\sum_{\alpha} f_{1, \alpha}^{\Sigma, P} \otimes \dots \otimes f_{v, \alpha}^{\Sigma, P} \in (U'^{\widehat{\otimes} n})^{\bar{\otimes} v}$.

Remark 4.4. One can prove that in the right hand side of (4.9), the contribution of all terms with $k = n$ is $(\text{Ad}(F^{(P)}) - \text{id}) \circ \delta^{(P)}$ where $F^{(P)}$ is the product of $F^{I,J}$ (I, J subsets of $\{1, \dots, n\}$, such that $\max(I) < \min(J)$) and their inverses such that

$$\Delta_F^{(P)} = \text{Ad}(F^{(P)}) \circ \Delta^{(P)}.$$

PROOF OF THE LEMMA. equation (4.9) may be proved by induction on $|P|$. Let us prove it for the unique tree P such that $|P| = 2$:

$$\delta_F^{(2)} = \delta^{(2)} + \sum_{v \geq 1} \frac{1}{v!} \text{ad}_{\bar{h}}(f)^v (\delta^{(2)}(x) + \delta^{(1)}(x)^1 + \delta^{(1)}(x)^2),$$

where (1) and (2) are the 1- and 2-trees. Assume that (4.9) is proved when $|P| = n$. Let P' be an $(n+1)$ -tree. Then for some $i \in \{1, \dots, n\}$, we have

$$\delta_F^{(P)} = (\text{id}^{\otimes i-1} \otimes \delta_F^{(2)} \otimes \text{id}^{\otimes n-i}) \circ \delta_F^{(P')},$$

where $|P'| = n$. Then:

$$\begin{aligned} \delta_F^{(P)} &= (\text{id}^{\otimes i-1} \otimes \Delta_F \otimes \text{id}^{\otimes n-i}) \circ \delta_F^{(P')} - (\delta_F^{(P')})_{1, \dots, \widehat{i}, \dots, n+1} - (\delta_F^{(P')})_{1, \dots, \widehat{i+1}, \dots, n+1} \\ &= (\text{id}^{\otimes i-1} \otimes \Delta_F \otimes \text{id}^{\otimes n-i}) \circ (\delta^{(P')} + \sum_{k \leq n} \sum_{R \text{ a } k\text{-tree}} \sum_{\substack{\Sigma \subset \{1, \dots, n\} \\ \text{card}(\Sigma)=k}} \\ &\quad \sum_{v \geq 1} \sum_{\alpha} \text{ad}_{\bar{h}}(f_{1, \alpha}^{\Sigma, P'}) \circ \dots \circ \text{ad}_{\bar{h}}(f_{v, \alpha}^{\Sigma, P'}) \circ (\delta^{(R)})^{\Sigma}) \\ &\quad - (\dots)_{1, \dots, \widehat{i}, \dots, n+1} - (\dots)_{1, \dots, \widehat{i+1}, \dots, n+1} \\ &= \text{Ad}(F^{i, i+1}) \circ (\delta^{(P)} + (\delta^{(P')})_{1, \dots, \widehat{i}, \dots, n+1} + (\delta^{(P')})_{1, \dots, \widehat{i+1}, \dots, n+1} \\ &\quad + \sum_{k \leq n} \sum_{R \text{ a } k\text{-tree}} \sum_{\substack{\Sigma \subset \{1, \dots, n\} \\ \text{card}(\Sigma)=k}} \sum_{v \geq 1} \sum_{\alpha} \text{ad}_{\bar{h}}((f_{1, \alpha}^{\Sigma, P'})_{1, \dots, \{i, i+1\}, \dots, n+1}) \circ \\ &\quad \circ \text{ad}_{\bar{h}}((f_{v, \alpha}^{\Sigma, P'})_{1, \dots, \{i, i+1\}, \dots, n+1}) \circ (1^{\otimes i-1} \otimes \Delta \otimes 1^{\otimes n-i}) \circ (\delta^{(R)})^{\Sigma}) \\ &\quad - (\dots)_{1, \dots, \widehat{i}, \dots, n+1} - (\dots)_{1, \dots, \widehat{i+1}, \dots, n+1}; \end{aligned}$$

this has the desired form because:

$$\begin{aligned} &(\text{Ad}(F^{i, i+1}) - 1) \circ (\delta^{(P)} + (\delta^{(P')})_{1, \dots, \widehat{i}, \dots, n+1} + (\delta^{(P')})_{1, \dots, \widehat{i+1}, \dots, n+1}) \\ &= \sum_{v \geq 1} \frac{1}{v!} \text{ad}_{\bar{h}}(f^{i, i+1})^v (\delta^{(P)} + (\delta^{(P')})_{1, \dots, \widehat{i}, \dots, n+1} + (\delta^{(P')})_{1, \dots, \widehat{i+1}, \dots, n+1}). \end{aligned}$$

This proves (4.9). $\square$

END OF PROOF OF PROPOSITION 4.2. One repeats the proof of Proposition 3.2 to prove that if $x \in U'$, then we have $\delta^{(P)}(x) \in \bar{h}^{|P|} U^{\otimes |P|}$ for any tree P . So $U' \subset U_F'$. Since (U, m, Δ, Φ) is the twist by F^{-1} of $(U, m, {}^F\Delta, {}^F\Phi)$, and $\bar{h} \log(F^{-1}) = -\bar{h} \log(F) \in (U')^{\otimes 2} \subset (U_F')^{\otimes 2}$, F^{-1} is admissible for $(U, m, {}^F\Delta, {}^F\Phi)$, so we have also $U_F' \subset U'$, so $U_F' = U'$. $\square$

- **b** - Twisting any algebra into an admissible algebra

Proposition 4.5. *Let (U, m, Δ, Φ) be a quasi-Hopf algebra. There exists a twist F_0 such that the twisted quasi-Hopf algebra $(U, m, {}^{F_0}\Delta, {}^{F_0}\Phi)$ is admissible.*

PROOF. We construct F_0 as a convergent infinite product $F_0 = \cdots F_n \cdots F_2$, where $F_n \in 1 + \hbar^{-1}U^{\widehat{\otimes} 2}$, and the F_n have the following property: if $\bar{F}_n = F_n F_{n-1} \cdots F_2$, if $\Phi_n = \bar{F}_n \Phi$, and $\delta_n^{(P)} : U \rightarrow U^{\widehat{\otimes} |P|}$ is the map corresponding to a tree P and to $\Delta_n = \text{Ad}(\bar{F}_n) \circ \Delta$, then we have

$$(\delta_n^{(P)} \otimes \delta_n^{(Q)} \otimes \delta_n^{(R)})(\hbar \log(\Phi_n)) \in \hbar^{|P|+|Q|+|R|} U^{\widehat{\otimes} |P|+|Q|+|R|}$$

for any trees P, Q, R such that $|P| + |Q| + |R| \leq n$.

Assume that we have constructed $F_1, \dots, F_n$, and let us construct F_{n+1} . The argument of Proposition 3.2 shows that for any integers (n_1, n_2, n_3) such that $n_1 + n_2 + n_3 = n + 1$, and any trees P, Q, R such that $|P| = n_1, |Q| = n_2, |R| = n_3$,

$$\left(\frac{1}{\hbar^n} (\delta_n^{(P)} \otimes \delta_n^{(Q)} \otimes \delta_n^{(R)})(\hbar \log(\Phi_n)) \bmod \hbar\right) \in S^{n_1}(\mathfrak{g}) \otimes S^{n_2}(\mathfrak{g}) \otimes S^{n_3}(\mathfrak{g}),$$

and is independent of the trees P, Q, R . The direct sum of these elements is an element $\bar{\varphi}_n$ of $S(\mathfrak{g})^{\otimes 3}$, homogeneous of degree $n + 1$. Since Φ_n satisfies the pentagon equation

$$(\text{id} \otimes \text{id} \otimes \Delta_n)(\Phi_n)^{-1} (1 \otimes \Phi_n) (\text{id} \otimes \Delta_n \otimes \text{id})(\Phi_n) (\Phi_n \otimes 1) (\Delta_n \otimes \text{id} \otimes \text{id})(\Phi_n)^{-1} = 1,$$

$\varphi_n^{\hbar} := \hbar \log(\Phi_n)$ satisfies the equation

$$\begin{aligned} & \left(-(\text{id} \otimes \text{id} \otimes \Delta_n)(\varphi_n^{\hbar}) \right) \star (1 \otimes \varphi_n^{\hbar}) \star \left((\text{id} \otimes \Delta_n \otimes \text{id})(\varphi_n^{\hbar}) \right) \star \\ & (\varphi_n^{\hbar} \otimes 1) \star \left(-(\Delta_n \otimes \text{id} \otimes \text{id})(\varphi_n^{\hbar}) \right) = 0, \end{aligned} \quad (4.10)$$

where we set

$$a \star b = a + b + \frac{1}{2}[a, b]_{\hbar} + \cdots$$

(the CBH series for the Lie bracket $[-, -]_{\hbar}$). Let (n_1, n_2, n_3, n_4) be integers such that $n_1 + \cdots + n_4 = n + 1$. Let P, Q, R, S be trees such that $|P| = n_1, \dots, |S| = n_4$. Let us apply $\delta_n^{(P)} \otimes \cdots \otimes \delta_n^{(S)}$ to (4.10). The left hand side of (4.10) is equal to

$$(-\Delta_n \otimes \text{id} \otimes \text{id} + \text{id} \otimes \Delta_n \otimes \text{id} - \text{id} \otimes \text{id} \otimes \Delta_n)(\varphi_n^{\hbar}) + (1 \otimes \varphi_n^{\hbar}) - (\varphi_n^{\hbar} \otimes 1) + \text{brackets}.$$

Now

$$(\delta_n^{(P)} \otimes \delta_n^{(Q)} \otimes \delta_n^{(R)} \otimes \delta_n^{(S)})(\Delta_n \otimes \text{id} \otimes \text{id})(\varphi_n^{\hbar}) = (\delta_n^{(P \cup Q)} \otimes \delta_n^{(R)} \otimes \delta_n^{(S)}).$$

where $P \cup Q$ is the tree with left descendant P and right descendant Q . Therefore

$$\left(\frac{1}{\hbar^n} (\delta_n^{(P)} \otimes \delta_n^{(Q)} \otimes \delta_n^{(R)} \otimes \delta_n^{(S)})(\Delta_n \otimes \text{id} \otimes \text{id})(\varphi_n^{\hbar}) \bmod \hbar\right) = (\Delta_0 \otimes \text{id} \otimes \text{id})(\bar{\varphi}_n)_{n_1, n_2, n_3, n_4}$$

where the index $(n_1, \dots, n_4)$ means the component in $\otimes_{i=1}^4 S^{n_i}(\mathfrak{g})$. On the other hand, if a_1 and $a_2 \in U^{\widehat{\otimes} 4}$ are such that

$$(\delta_n^{(P)} \otimes \cdots \otimes \delta_n^{(S)})(a_i) \in \hbar^{\inf(|P| + \cdots + |S|, n)} U^{\widehat{\otimes} 4}$$

for any trees $(P, \dots, S)$, then if $(P, \dots, S)$ are such that $|P| + \cdots + |S| = n$, we have

$$(\delta_n^{(P)} \otimes \cdots \otimes \delta_n^{(S)})(\frac{1}{\hbar}[a_1, a_2]) \in \hbar^{n+1} U^{\widehat{\otimes} n};$$

one proves this in the same way as the commutativity of $U'/\hbar U'$ (see Theorem 1.1). Then $\frac{1}{\hbar^n}(\delta_n^{(P)} \otimes \cdots \otimes \delta_n^{(S)})(4.10)|_{\hbar=0}$ yields $d(\bar{\varphi}_n) = 0$, where $d : S(\mathfrak{g})^{\otimes 2} \rightarrow S(\mathfrak{g})^{\otimes 3}$ is the co-Hochschild cohomology differential. This relation implies that

$$\bar{\varphi}_n = d(\bar{f}_n) + \lambda_n,$$

where $\bar{f}_n \in S(\mathfrak{g})^{\otimes 2}$ and $\lambda_n \in \Lambda^3(\mathfrak{g})$. Moreover, f_n and λ_n both have degree $n+1$. This implies that $\lambda_n = 0$. Let $f_n \in (U(\mathfrak{g})^{\otimes 2})_{\leq n+1}$ be a preimage of $\bar{f}_n$ by the projection

$$(U(\mathfrak{g})^{\otimes 2})_{\leq n+1} \rightarrow (U(\mathfrak{g})^{\otimes 2})_{\leq n+1} / (U(\mathfrak{g})^{\otimes 2})_{\leq n} = (S(\mathfrak{g})^{\otimes 2})_{n+1}$$

(where the indices n and $\leq n$ mean “homogeneous part of degree n ” and “part of degree $\leq n$ ”). Let $f_n^{\hbar} \in U^{\hat{\otimes} 2}$ be a preimage of f_n by the projection $U^{\hat{\otimes} 2} \rightarrow U^{\hat{\otimes} 2} / \hbar U^{\hat{\otimes} 2} = U(\mathfrak{g})^{\otimes 2}$. Set $F_{n+1} = \exp(\hbar^{n-1} f_n)$. We may assume that $\hbar^n f_n \in (U(\bar{F}_n)')^{\otimes 2}$, where $U(\bar{F}_n)' = \{x \in U | \delta_n^{(P)}(x) \in \hbar^{\inf(n, |P|)} U^{\hat{\otimes} |P|}\}$. Then $\Phi_{n+1} = F_{n+1} \Phi_n$. If P, Q, R are such that $|P| + |Q| + |R| = n+1$, then

$$(\delta_n^{(P)} \otimes \delta_n^{(Q)} \otimes \delta_n^{(R)})(\hbar \log(\Phi_{n+1})) \in \hbar^{n+1} U^{\hat{\otimes} n+1}.$$

Then according to Lemma 4.3,

$$(\delta_{n+1}^{(P)} \otimes \delta_{n+1}^{(Q)} \otimes \delta_{n+1}^{(R)} - \delta_n^{(P)} \otimes \delta_n^{(Q)} \otimes \delta_n^{(R)})(\hbar \log(\Phi_{n+1}))$$

has $\hbar$ -adic valuation $> |P| + |Q| + |R|$ when $|P| + |Q| + |R| \leq n+1$. So $(\delta_{n+1}^{(P)} \otimes \delta_{n+1}^{(Q)} \otimes \delta_{n+1}^{(R)})(\hbar \log(\Phi_{n+1})) \in \hbar^{|P|+|Q|+|R|} U^{\hat{\otimes} |P|+|Q|+|R|}$ whenever $|P| + |Q| + |R| \leq n+1$. $\square$

§ 5 Proof of Proposition 2.4

We work by induction on n . The statement is obvious when $n = 0, 1$. For $n = 2$, we get

$$\begin{aligned} \delta^{(2)}(xy) &= \delta^{(2)}(x)\delta^{(2)}(y) + \delta^{(2)}(x)(\delta^{(1)}(y)^1 + \delta^{(1)}(y)^2 + \delta^{(0)}(y)^0) \\ &\quad + (\delta^{(1)}(x)^1 + \delta^{(1)}(y)^1 + \delta^{(0)}(y)^0)\delta^{(2)}(y) \\ &\quad + \delta^{(1)}(x)^1\delta^{(2)}(y)^2 + \delta^{(1)}(x)^2\delta^{(2)}(y)^1, \end{aligned} \tag{5.11}$$

so the statement also holds.

Assume that the statement is proved when P is a n -tree. Let $\bar{P}$ be a $(n+1)$ -tree. There exists an integer $k \in \{0, \dots, n-1\}$, such that $\bar{P}$ may be viewed as the glueing of the 2-tree on the k -th leaf of a n -tree P . Then we have

$$\delta^{(\bar{P})} = (\text{id}^{\otimes k} \otimes \delta^{(2)} \otimes \text{id}^{\otimes n-k-1}) \circ \delta^{(P)}.$$

Let us assume, for instance, that $k = n-1$. If v is an integer, set

$$S_v = \{(\Sigma', \Sigma'') | \Sigma', \Sigma'' \subset \{1, \dots, v\} \text{ and } \Sigma' \cup \Sigma'' = \{1, \dots, v\}\}.$$

Then

$$S_n = f_{\{n\}, \emptyset}(S_{n-1}) \cup f_{\emptyset, \{n\}}(S_{n-1}) \cup f_{\{n\}, \{n\}}(S_{n-1}) \text{ (disjoint union),}$$

where $f_{\alpha, \beta}(\Sigma', \Sigma'') = (\Sigma' \cup \alpha, \Sigma'' \cup \beta)$. By hypothesis, we have

$$\delta^{(P)}(xy) = \sum_{(\Sigma_1, \Sigma_2) \in S_n} \delta^{(P_{\Sigma_1})}(x)^{\Sigma_1} \delta^{(P_{\Sigma_2})}(y)^{\Sigma_2},$$

therefore

$$\begin{aligned}\delta^{(P)}(xy) &= \sum_{(\Sigma', \Sigma'') \in S_{n-1}} \delta^{(P_{\Sigma' \cup \{n\}})}(x)^{\Sigma' \cup \{n\}} \delta^{(P_{\Sigma''})}(y)^{\Sigma''} \\ &\quad + \delta^{(P_{\Sigma'})}(x)^{\Sigma'} \delta^{(P_{\Sigma'' \cup \{n\}})}(y)^{\Sigma'' \cup \{n\}} \\ &\quad + \delta^{(P_{\Sigma' \cup \{n\}})}(x)^{\Sigma' \cup \{n\}} \delta^{(P_{\Sigma'' \cup \{n\}})}(y)^{\Sigma'' \cup \{n\}}.\end{aligned}$$

Applying $\text{id}^{\otimes n-1} \otimes \delta^{(2)}$ to this identity and using (5.11) and the identities

$$\begin{aligned}(\text{id}^{\otimes k} \otimes \delta^{(1)} \otimes \text{id}^{\otimes |P|-k-1}) \circ \delta^{(P)} &= \delta^{(P)}, \\ (\text{id}^{\otimes k} \otimes \delta^{(0)} \otimes \text{id}^{\otimes |P|-k-1}) \circ \delta^{(P)} &= 0,\end{aligned}$$

we get $\delta^{(\bar{P})}(xy) =$

$$\begin{aligned}&\sum_{(\Sigma', \Sigma'') \in S_{n-1}} \left((\text{id}^{\otimes |\Sigma'|} \otimes \delta^{(2)}) \circ \delta^{(P_{\Sigma' \cup \{n\}})}(x)^{\Sigma' \cup \{n, n+1\}} \delta^{(P_{\Sigma''})}(y)^{\Sigma''} \right. \\ &\quad + \delta^{(P_{\Sigma'})}(x)^{\Sigma'} \left((\text{id}^{\otimes |\Sigma''|} \otimes \delta^{(2)}) \circ \delta^{(P_{\Sigma'' \cup \{n\}})}(y)^{\Sigma'' \cup \{n, n+1\}} \right. \\ &\quad + \left((\text{id}^{\otimes |\Sigma'|} \otimes \delta^{(2)}) \circ \delta^{(P_{\Sigma' \cup \{n\}})}(x)^{\Sigma' \cup \{n, n+1\}} \right) \left((\text{id}^{\otimes |\Sigma''|} \otimes \delta^{(2)}) \circ \delta^{(P_{\Sigma'' \cup \{n\}})}(y)^{\Sigma'' \cup \{n, n+1\}} \right. \\ &\quad + \left((\text{id}^{\otimes |\Sigma'|} \otimes \delta^{(2)}) \circ \delta^{(P_{\Sigma' \cup \{n\}})}(x)^{\Sigma' \cup \{n, n+1\}} \right) \left(\delta^{(P_{\Sigma'' \cup \{n\}})}(y)^{\Sigma'' \cup \{n\}} + \delta^{(P_{\Sigma'' \cup \{n+1\}})}(y)^{\Sigma'' \cup \{n+1\}} \right) \\ &\quad + \left(\delta^{(P_{\Sigma' \cup \{n\}})}(x)^{\Sigma' \cup \{n\}} + \delta^{(P_{\Sigma' \cup \{n+1\}})}(x)^{\Sigma' \cup \{n+1\}} \right) \left((\text{id}^{\otimes |\Sigma''|} \otimes \delta^{(2)}) \circ \delta^{(P_{\Sigma'' \cup \{n\}})}(y)^{\Sigma'' \cup \{n, n+1\}} \right. \\ &\quad \left. \left. + \delta^{(P_{\Sigma' \cup \{n\}})}(x)^{\Sigma' \cup \{n\}} \delta^{(P_{\Sigma'' \cup \{n\}})}(y)^{\Sigma'' \cup \{n+1\}} + \delta^{(P_{\Sigma' \cup \{n+1\}})}(x)^{\Sigma' \cup \{n+1\}} \delta^{(P_{\Sigma'' \cup \{n\}})}(y)^{\Sigma'' \cup \{n\}} \right) \right).\end{aligned}$$

So we get $\delta^{(\bar{P})}(xy) =$

$$\begin{aligned}&\sum_{(\Sigma', \Sigma'') \in S_{n-1}} \left(\delta^{(\bar{P}_{\Sigma' \cup \{n, n+1\}})}(x)^{\Sigma' \cup \{n, n+1\}} \delta^{(\bar{P}_{\Sigma''})}(y)^{\Sigma''} \right. \\ &\quad + \delta^{(\bar{P}_{\Sigma'})}(x)^{\Sigma'} \delta^{(\bar{P}_{\Sigma'' \cup \{n, n+1\}})}(y)^{\Sigma'' \cup \{n, n+1\}} \\ &\quad + \delta^{(\bar{P}_{\Sigma' \cup \{n, n+1\}})}(x)^{\Sigma' \cup \{n, n+1\}} \delta^{(\bar{P}_{\Sigma'' \cup \{n, n+1\}})}(y)^{\Sigma'' \cup \{n, n+1\}} \\ &\quad + \delta^{(\bar{P}_{\Sigma' \cup \{n, n+1\}})}(x)^{\Sigma' \cup \{n, n+1\}} \left(\delta^{(\bar{P}_{\Sigma'' \cup \{n\}})}(y)^{\Sigma'' \cup \{n\}} + \delta^{(\bar{P}_{\Sigma'' \cup \{n+1\}})}(y)^{\Sigma'' \cup \{n+1\}} \right) \\ &\quad + \left(\delta^{(\bar{P}_{\Sigma' \cup \{n\}})}(x)^{\Sigma' \cup \{n\}} + \delta^{(\bar{P}_{\Sigma' \cup \{n+1\}})}(x)^{\Sigma' \cup \{n+1\}} \right) \delta^{(\bar{P}_{\Sigma'' \cup \{n, n+1\}})}(y)^{\Sigma'' \cup \{n, n+1\}} \\ &\quad \left. + \delta^{(\bar{P}_{\Sigma' \cup \{n\}})}(x)^{\Sigma' \cup \{n\}} \delta^{(\bar{P}_{\Sigma'' \cup \{n+1\}})}(y)^{\Sigma'' \cup \{n+1\}} + \delta^{(\bar{P}_{\Sigma' \cup \{n+1\}})}(x)^{\Sigma' \cup \{n+1\}} \delta^{(\bar{P}_{\Sigma'' \cup \{n\}})}(y)^{\Sigma'' \cup \{n\}} \right).\end{aligned}$$

We have

$$\begin{aligned}S_{n+1} &= f_{\{n, n+1\}, \{n, n+1\}}(S_{n-1}) \cup f_{\{n, n+1\}, \{n\}}(S_{n-1}) \cup f_{\{n, n+1\}, \{n+1\}}(S_{n-1}) \\ &\quad \cup f_{\{n, n+1\}, \emptyset}(S_{n-1}) \cup f_{\{n\}, \{n, n+1\}}(S_{n-1}) \cup f_{\{n+1\}, \{n, n+1\}}(S_{n-1}) \\ &\quad \cup f_{\emptyset, \{n, n+1\}}(S_{n-1}) \cup f_{\{n\}, \{n+1\}}(S_{n-1}) \cup f_{\{n+1\}, \{n\}}(S_{n-1}) \text{ (disjoint union),}\end{aligned}$$

where we recall that $f_{\alpha, \beta}(\Sigma', \Sigma'') = (\Sigma' \cup \alpha, \Sigma'' \cup \beta)$. So we get

$$\delta^{(\bar{P})}(xy) = \sum_{(\bar{\Sigma}', \bar{\Sigma}'') \in S_{n+1}} \delta^{(P_{\bar{\Sigma}'})}(x)^{|\bar{\Sigma}'|} \delta^{(P_{\bar{\Sigma}''})}(y)^{|\bar{\Sigma}''|}.$$

The proof is the same for a general $k \in \{0, \dots, n-1\}$. This establishes the induction. $\square$

§ 6 Proofs of Proposition 1.4, Theorem 1.8 and Proposition 1.9

1. *Proof of Proposition 1.4.* According to [Dr2], Proposition 3.10, there exists a series $\mathcal{E}'(\varphi) \in U(\mathfrak{g})^{\otimes 3}[[\hbar]]$, expressed in terms of (μ, φ) by universal acyclic expressions (and therefore invariant), such that $\mathcal{E}'(\varphi) = 1 + O(\hbar^2)$, and $\mathcal{E}'(\varphi)$ satisfies the pentagon identity. Then $(U(\mathfrak{g})[[\hbar]], m_0, \Delta_0, \mathcal{E}'(\varphi))$ is a quasi-Hopf algebra. By Theorem 1.3, 2), there exists a twist $F \in U(\mathfrak{g})^{\otimes 2}[[\hbar]]^\times$, such that $(U(\mathfrak{g})[[\hbar]], m_0, {}^F\Delta_0, {}^F\mathcal{E}'(\varphi))$ is admissible.

$\mathcal{E}(\varphi)$ gives rise to a collection of invariant elements $\mathcal{E}'(\varphi)_{p_1, p_2, p_3, n} \in \otimes_{i=1}^3 S^{p_i}(\mathfrak{g})$, defined by the condition that the image of $\mathcal{E}'(\varphi)$ by the symmetrization map $U(\mathfrak{g})^{\otimes 3}[[\hbar]] \rightarrow S(\mathfrak{g})[[\hbar]]$ is $\sum_{n \geq 0, p_1, p_2, p_3 \geq 0} \hbar^n \mathcal{E}'(\varphi)_{p_1, p_2, p_3, n}$. F is then expressed using only the $\mathcal{E}'_{p_1, p_2, p_3, n}$, the Lie bracket and the symmetric group operations on the $\mathfrak{g}^{\otimes n}$. So F is invariant and defined by universal acyclic expressions. Therefore ${}^F\Delta_0 = \Delta_0$. $\mathcal{E}(\varphi) := {}^F\mathcal{E}'(\varphi)$ is then expressed by universal acyclic expressions, and defines an admissible quantization of $(\mathfrak{g}, \mu, \delta = 0, \varphi)$.

2. *Proof of Theorem 1.8, 1).* We have then $\mathcal{E}(\varphi) \in (U(\mathfrak{g})[[\hbar]]')^{\bar{\otimes} 3}$. Since the coproduct is Δ_0 , $U(\mathfrak{g})[[\hbar]]'$ is the complete subalgebra of $U(\mathfrak{g})[[\hbar]]$ generated by $\hbar\mathfrak{g}$, so it is a flat deformation of $\hat{S}(\mathfrak{g})$ with Kostant-Kirillov Poisson structure. We then set $\tilde{\varphi} := \mathcal{E}(\varphi)$ modulo $\hbar$.

3. *Proof of Theorem 1.8, 2).* Let $\tilde{\varphi}_1, \tilde{\varphi}_2$ be the elements of $\hat{S}(\mathfrak{g})^{\bar{\otimes} 3}$ such that

$$(\hat{S}(\mathfrak{g}), m_0, P_{\mathfrak{g}}, \Delta_0, \tilde{\varphi}_i)$$

are Drinfeld algebras. Let C be the lowest degree component of $\tilde{\varphi}_1 - \tilde{\varphi}_2$. Then the degree k of C is ≥ 4 . Taking the degree k part of the difference of the pentagon identities for $\tilde{\varphi}_1$ and $\tilde{\varphi}_2$, we find $d(C) = 0$, where $d : S(\mathfrak{g})^{\otimes 3} \rightarrow S(\mathfrak{g})^{\otimes 4}$ is the co-Hochschild differential. So $\text{Alt}(C) \in \Lambda^3(\mathfrak{g})$, and since $\text{Alt}(C)$ also has degree ≥ 4 , $\text{Alt}(C) = 0$. If C_{p_1, p_2, p_3} is the component of C in $\otimes_{i=1}^3 S^{p_i}(\mathfrak{g})$ then we may define inductively B by $B_{0, k} = B_{1, k-1} = 0$, $B_{2, k-2} = \frac{1}{2}(\text{id} \otimes m)(C_{1, 1, k-2})$, and

$$B_{i+1, k-i-1} = \frac{1}{i+1}(\text{id} \otimes m)[C_{i, 1, k-i-1} + ((\text{id} \otimes d)(B_{i, k-i}))_{i, 1, k-i-1}],$$

where $B_{i, j}$ is the component of B in $S^i(\mathfrak{g}) \otimes S^j(\mathfrak{g})$ and m is the product of $S(\mathfrak{g})$. So B can be chosen to be $\mathfrak{g}$ -invariant. Applying successive twists, we obtain the result.

4. *Proposition 1.9.* According to [Dr3], $(U(\mathfrak{g}), m_0, \Delta_0, e^{\hbar\mu_{\mathfrak{g}}/2}, \Phi(\hbar t_{\mathfrak{g}}^{1,2}, \hbar t_{\mathfrak{g}}^{2,3}))$ is a quasi-triangular quasi-Hopf algebra. One checks that it is admissible; then the reduction modulo $\hbar$ of the corresponding QFS algebra is the Drinfeld algebra of 1). $\square$

Remark 6.1. In the proof of Theorem 1.8, 2), we cannot use Theorem A of [Dr2] because we do not know that the twist constructed there is admissible.

§ 7 Associators and Lie associators

In this section, we state precisely and prove Theorem 1.10.

- a - Statement of the result

Recall that the algebra $\mathcal{T}_n$, $n \geq 2$, has generators $t^{i,j}$, $1 \leq i \neq j \leq n$, and relations $t^{j,i} = t^{i,j}$,

$$\begin{aligned} [t^{i,j} + t^{i,k}, t^{j,k}] &= 0 \text{ when } i, j, k \text{ are all distinct,} \\ [t^{i,j}, t^{k,l}] &= 0 \text{ when } i, j, k, l \text{ are all distinct.} \end{aligned}$$

$\mathfrak{t}_n$ is defined as the Lie algebra with the same generators and relations. Then $\mathcal{T}_n = U(\mathfrak{t}_n)$. When $n \leq m$ and $(I_1, \dots, I_n)$ is a collection of disjoint subsets of $\{1, \dots, m\}$, there is a unique algebra morphism $\mathcal{T}_n \rightarrow \mathcal{T}_m$ taking $t^{i,j}$ to $\sum_{\alpha \in I_i, \beta \in I_j} t^{\alpha, \beta}$. We call it an insertion-coproduct morphism and denote it by $x \mapsto x^{I_1, \dots, I_n}$. In particular, we have an action of $\mathfrak{S}_n$ on $\mathcal{T}_n$. Let us attribute degree 1 to each generator $t^{i,j}$; this defines gradings on the algebra $\mathcal{T}_n$ and on the Lie algebra $\mathfrak{t}_n$. We denote by $\widehat{\mathcal{T}}_n$ and $\widehat{\mathfrak{t}}_n$ their completions for this grading. Then $\widehat{\mathcal{T}}_n$ is the preimage of $\mathbb{K}^\times$ by the natural projection $\widehat{\mathcal{T}}_n \rightarrow \mathbb{K}$, and the exponential is a bijection $(\widehat{\mathcal{T}}_n)_0 \rightarrow 1 + (\widehat{\mathcal{T}}_n)_0$ (where $(\widehat{\mathcal{T}}_n)_0 = \text{Ker}(\widehat{\mathcal{T}}_n \rightarrow \mathbb{K})$). We have an exact sequence

$$1 \rightarrow 1 + (\widehat{\mathcal{T}}_n)_0 \rightarrow (\widehat{\mathcal{T}}_n)^\times \rightarrow \mathbb{K}^\times \rightarrow 1.$$

An *associator* is an element Φ of $1 + (\widehat{\mathcal{T}}_n)_0$, satisfying the pentagon equation

$$\Phi^{1,2,34} \Phi^{12,3,4} = \Phi^{2,3,4} \Phi^{1,23,4} \Phi^{1,2,3}, \quad (7.12)$$

the hexagon equations

$$e^{\frac{t^{1,3} + t^{2,3}}{2}} = \Phi^{3,1,2} e^{\frac{t^{1,3}}{2}} (\Phi^{1,3,2})^{-1} e^{\frac{t^{2,3}}{2}} \Phi^{1,2,3}$$

and

$$e^{\frac{t^{1,2} + t^{1,3}}{2}} = (\Phi^{2,3,1})^{-1} e^{\frac{t^{1,3}}{2}} \Phi^{2,1,3} e^{\frac{t^{1,2}}{2}} (\Phi^{1,2,3})^{-1}$$

and $\text{Alt}(\Phi) = \frac{1}{8}[t^{1,2}, t^{2,3}] + \text{terms of degree } > 2$. We denote by Assoc the set of associators. If Φ satisfies the duality condition $\Phi^{3,2,1} = \Phi^{-1}$, then both hexagon equations are equivalent. We denote by Assoc⁰ the subset of all $\Phi \in \text{Assoc}$ satisfying the duality condition. If $F \in 1 + (\widehat{\mathcal{T}}_2)_0$ and $\Phi \in 1 + (\widehat{\mathcal{T}}_3)_0$, the *twist of Φ by F* is

$${}^F\Phi = F^{2,3} F^{1,23} \Phi (F^{1,2} F^{12,3})^{-1}.$$

This defines an action of $1 + (\widehat{\mathcal{T}}_2)_0$ on $1 + (\widehat{\mathcal{T}}_3)_0$, which preserves Pent = $\{\Phi \in 1 + (\widehat{\mathcal{T}}_3)_0 \mid \Phi \text{ satisfies (7.12)}\}$, Assoc and Assoc⁰ (Pent and Assoc are preserved because F has the form $f(t^{1,2})$, $f \in 1 + t\mathbb{K}[[t]]$, so the “twisted R -matrix” ${}^F R = F^{2,1} R F^{-1} = f(t^{2,1}) e^{t^{1,2}/2} f(t^{1,2})^{-1} = e^{t^{1,2}/2}$. Assoc⁰ is preserved because each F is such that $F = F^{2,1}$.) We denote by Assoc_{Lie}⁰, Assoc_{Lie} and Pent_{Lie} the subsets of all Φ in Assoc, Assoc⁰ and Pent, such that $\log(\Phi) \in \widehat{\mathfrak{t}}_3$.

Theorem 7.1. *There is exactly one element of Pent_{Lie} resp., Assoc_{Lie}, Assoc_{Lie}⁰ in each orbit of the action of $1 + (\widehat{\mathcal{T}}_2)_0$ on Pent (resp., Assoc, Assoc⁰). The isotropy group of each element of Pent is $\{e^{\lambda t^{1,2}} \mid \lambda \in \mathbb{K}\} \subset 1 + (\widehat{\mathcal{T}}_2)_0$.*

- b - Proof of Theorem 7.1

The arguments are the same in all three cases, so we treat the case of Assoc.

Let Φ belongs to Assoc. Set $\Phi = 1 + \sum_{i>0} \Phi_i$, where Φ_i is the degree i component of Φ .

Let d be the co-Hochschild differential,

$$d : \mathcal{T}_n \rightarrow \mathcal{T}_{n+1}$$

$$x \mapsto \sum_{i=1}^n (-1)^{i+1} x^{1,\dots,\{i,i+1\},\dots,n+1} - x^{2,3,\dots,n+1} + (-1)^n x^{1,2,\dots,n}.$$

Then $d(\Phi_2) = 0$, and $\text{Alt}(\Phi_2) = \frac{1}{8}[t^{1,2}, t^{2,3}]$. Computation shows that this implies that for some $\lambda \in \mathbb{K}$, we have $\Phi_2 = \frac{1}{8}[t^{1,2}, t^{2,3}] + \lambda d((t^{1,2})^2)$. We construct $F \in 1 + (\widehat{\mathcal{T}}_2)_0$, such that ${}^F\Phi \in \underline{\text{Assoc}}_{\text{Lie}}$, as an infinite product $F = \cdots F_n \cdots F_2$, where $F_i \in 1 + (\widehat{\mathcal{T}}_2)_{\geq i}$ (the index $\geq i$ means the part of degree $\geq i$). If we set $F_2 = 1 + \lambda (t^{1,2})^2$, then $\log({}^F\Phi) \in \mathfrak{t}_3 + (\widehat{\mathcal{T}}_3)_{\geq 3}$. Assume that we have found $F_3, \dots, F_{n-1}$, such that $\log({}^{F_{n-1}}\Phi) \in \mathfrak{t}_3 + (\widehat{\mathcal{T}}_3)_{\geq n}$, where $F_{n-1} = F_{n-1} \cdots F_2$. Then $\varphi^{(n-1)} := \log({}^{F_{n-1}}\Phi)$ satisfies

$$(\varphi^{(n-1)})^{1,2,3,4} \star (\varphi^{(n-1)})^{12,3,4} = (\varphi^{(n-1)})^{2,3,4} \star (\varphi^{(n-1)})^{1,23,4} \star (\varphi^{(n-1)})^{1,2,3},$$

where $\star$ is the CBH product in $(\widehat{\mathcal{T}}_3)_0$. Let $\varphi_n^{(n-1)}$ be the degree n part of $\varphi^{(n-1)}$. Then we get $d(\varphi^{(n-1)}) \in \mathfrak{t}_4$. We now use the following statement, which will be proved in the next subsection.

Proposition 7.2. *If $\gamma \in \mathcal{T}_3$ is such that $d(\gamma) \in \mathfrak{t}_4$, then there exists $\beta \in \mathcal{T}_2$, such that $\gamma + d(\beta) \in \mathfrak{t}_3$. If γ has degree n , one can choose β of degree n .*

It follows that there exists $\beta \in \mathcal{T}_2$ of degree n , such that $\varphi_n^{(n-1)} - d(\beta) \in \mathfrak{t}_3$. Set $F_n = 1 + \beta$, then $\varphi^{(n)} = \log({}^{F_n}\Phi)$ is such that $\varphi^{(n)} \in \varphi^{(n-1)} - d(\beta) + (\widehat{\mathcal{T}}_3)_{\geq n+1}$, so $\varphi^{(n)} \in \mathfrak{t}_3 + (\widehat{\mathcal{T}}_3)_{\geq n+1}$. Moreover, the product $F = \cdots F_n \cdots F_2$ is convergent, and ${}^F\Phi$ then satisfies $\log({}^F\Phi) \in \widehat{\mathfrak{t}}_3$. This proves the existence of F , such that ${}^F\Phi \in \underline{\text{Assoc}}_{\text{Lie}}$.

Let us now prove the unicity of an element of Assoc_{Lie}, twist-equivalent to $\Phi \in \underline{\text{Assoc}}$. This follows from:

Proposition 7.3. *Let Φ' and Φ'' be elements of Assoc_{Lie}, and let F belong to $1 + (\widehat{\mathcal{T}}_2)_0$. Then ${}^F\Phi' = \Phi''$ if and only if there exists $\lambda \in \mathbb{K}$ such that $F = e^{\lambda t^{1,2}}$ and $\Phi' = \Phi''$.*

PROOF OF PROPOSITION 7.3. Since $t^{1,2} + t^{1,3} + t^{2,3}$ is central in $\widehat{\mathcal{T}}_3$, we have ${}^{F_\lambda}\Phi' = \Phi'$ when $F_\lambda = e^{\lambda t}$, for any $\lambda \in \mathbb{K}$. Conversely, let F_i be the degree i part of F . Then for some $\lambda_0 \in \mathbb{K}$, we have $F_1 = \lambda_0 t$. Replacing F by $F' = FF_{-\lambda_0}$, we get ${}^{F'}\Phi' = \Phi''$, and $F' - 1$ has valuation ≥ 2 (for the degree in t). Assume that $F' - 1 \neq 0$ and let v be its valuation. Let F'_v be the degree v part of F' . Then $d(F'_v) \in \mathfrak{t}_3$. On the other hand, $F'_v = \mu (t^{1,2})^v$, where $\mu \in \mathbb{K} - \{0\}$. Now $d((t^{1,2})^v) \in \mathcal{T}_3 = U(\mathfrak{t}_3)$ has degree $\leq v$ for the filtration of $U(\mathfrak{t}_3)$, and its symbol in $S^v(\mathfrak{t}_3) = \text{gr}_v(U(\mathfrak{t}_3))$ is $\sum_{v'=1}^{v-1} \binom{v}{v'} (t^{1,3})^{v'} (t^{2,3})^{v-v'} - \sum_{v''=1}^{v-1} \binom{v}{v''} (t^{1,2})^{v''} (t^{1,3})^{v-v''}$: this is the image of a non-zero element in $S^v(\mathbb{K}t^{1,2} \oplus \mathbb{K}t^{1,3} \oplus \mathbb{K}t^{2,3})$ under the injection $S^v(\bigoplus_{1 \leq i < j \leq 3} \mathbb{K}t^{i,j}) \hookrightarrow S^v(\mathfrak{t}_3)$, so it is non-zero. So $F' \neq 1$ leads to a contradiction. So $F = F_{\lambda_0}$, therefore $\Phi'' = \Phi'$. $\square$

Note that we have proved the analogue of Proposition 7.2, where the indices of $\mathcal{T}_3, \mathfrak{t}_4$, etc., are shifted by -1 .

- c - Decomposition of $\mathfrak{t}_3$ and proof of Proposition 7.2

To end the proof of the first part of Theorem 7.1, it remains to prove Proposition 7.2. For this, we construct a decomposition of $\mathfrak{t}_n$. For $i = 1, \dots, n$, there is a unique algebra morphism $\varepsilon_i : \mathcal{T}_n \rightarrow \mathcal{T}_{n-1}$, taking $t_{i,j}$ to 0 for any $j \neq i$, and taking $t_{j,k}$ to $t_{j-\lambda_i(j), k-\lambda_i(k)}$ if $j, k \neq i$, where $\lambda_i(j) = 0$ if $j < i$ and $= 1$ if $j > i$. Then ε_i induces a Lie algebra morphism $\tilde{\varepsilon}_i : \mathfrak{t}_n \rightarrow \mathfrak{t}_{n-1}$. Set $\tilde{\mathfrak{t}}_n = \bigcap_{i=1}^n \text{Ker}(\tilde{\varepsilon}_i)$. Then we have

Lemma 7.4.

$$\mathfrak{t}_n = \bigoplus_{k=0}^n \bigoplus_{I \in \mathcal{P}_k(\{1, \dots, n\})} (\tilde{\mathfrak{t}}_k)^I,$$

where $\mathcal{P}_k(\{1, \dots, n\})$ is the set of subsets of $\{1, \dots, n\}$ of cardinal k , and $(\tilde{\mathfrak{t}}_k)^I$ is the image of $\tilde{\mathfrak{t}}_k$ under $\mathfrak{t}_k \rightarrow \mathfrak{t}_n, x \mapsto x^{i_1 \dots i_k}$, where $I = \{i_1, \dots, i_k\}$.

PROOF OF LEMMA. Let $\mathfrak{F}$ be the free Lie algebra with generators $\tilde{t}_{i,j}$, where $1 \leq i < j \leq n$. It is graded by $\Gamma := \mathbb{N}^{\{(i,j) | 1 \leq i < j \leq n\}}$: the degree of $\tilde{t}_{i,j}$ is the vector $\mathbf{d}_{i,j}$, whose (i', j') coordinate is $\delta_{(i,j), (i', j')}$. For $\underline{k} \in \Gamma$, we denote by $\mathfrak{F}_{\underline{k}}$ the part of $\mathfrak{F}$ of degree $\underline{k}$. Let $\pi : \mathfrak{F} \rightarrow \mathfrak{t}_n$ be the canonical projection. Since the defining ideal of $\mathfrak{t}_n$ is graded, we have

$$\mathfrak{t}_n = \bigoplus_{\underline{k} \in \Gamma} \pi(\mathfrak{F}_{\underline{k}}). \quad (7.13)$$

On the other hand, one checks that $\tilde{\mathfrak{t}}_n = \bigoplus_{\underline{k} \in \tilde{\Gamma}} \pi(\mathfrak{F}_{\underline{k}})$, where $\tilde{\Gamma}$ is the set of maps $k : \{(i, j) | 1 \leq i < j \leq n\} \rightarrow \mathbb{N}$, such that for each i , $\sum_{j|j>i} k(i, j) + \sum_{j|j<i} k(j, i) \neq 0$. Define a map $\lambda : \Gamma \rightarrow \mathcal{P}(\{1, \dots, n\})$ as follows ($\mathcal{P}(\{1, \dots, n\})$ is the set of subsets of $\{1, \dots, n\}$): λ takes the map $k : \{(i, j) | 1 \leq i < j \leq n\} \rightarrow \mathbb{N}$ to $\{i | \sum_{j|j>i} k(i, j) + \sum_{j|j<i} k(j, i) \neq 0\}$. Then for each $I \in \mathcal{P}(\{1, \dots, n\})$, $(\tilde{\mathfrak{t}}_I)^I$ identifies with $\bigoplus_{\underline{k} \in \lambda^{-1}(I)} \pi(\mathfrak{F}_{\underline{k}})$. Comparing with (7.13), we get

$$\mathfrak{t}_n = \bigoplus_{I \in \mathcal{P}(\{1, \dots, n\})} (\tilde{\mathfrak{t}}_I)^I. \quad \square$$

When $n = 3$, we get $\mathfrak{t}_3 = \mathbb{K}t^{1,2} \oplus \mathbb{K}t^{1,3} \oplus \mathbb{K}t^{2,3} \oplus \tilde{\mathfrak{t}}_3$. On the other hand, the fact that the insertion-coproduct maps take $\mathfrak{t}_n$ to $\mathfrak{t}_m$ implies that $d : \mathcal{T}_n \rightarrow \mathcal{T}_{n+1}$ is compatible with the filtrations induced by the identification $\mathcal{T}_n = U(\mathfrak{t}_n)$, $\mathcal{T}_{n+1} = U(\mathfrak{t}_{n+1})$. The associated graded map is

$$\text{gr}(d) : S(\mathfrak{t}_n) \rightarrow S(\mathfrak{t}_{n+1}).$$

Proposition 7.2 now follows from:

Lemma 7.5. When $k \geq 2$, the cohomology of the complex

$$S^k(\mathfrak{t}_2) \xrightarrow{\text{gr}^k(d)} S^k(\mathfrak{t}_3) \xrightarrow{\text{gr}^k(d)} S^k(\mathfrak{t}_4)$$

vanishes.

PROOF OF LEMMA. We have

$$S^k(t_3) = \bigoplus_{\alpha=0}^k S^{k-\alpha} \left(\bigoplus_{1 \leq i < j \leq 3} \mathbb{K} t^{i,j} \right) \otimes S^\alpha(\tilde{t}_3). \quad (7.14)$$

Let $x \in S^k(t_3)$, and let $(x_\alpha)_{\alpha=0,\dots,k}$ be its components in the decomposition (7.14). We have

$$S(t_4) = S(\tilde{t}_4) \otimes \bigotimes_{2 \leq i < j \leq 4} S(\tilde{t}_3^{1,i,j}) \otimes \bigotimes_{i=2}^4 S(\tilde{t}_2^{1,i}) \otimes S(t_3^{2,3,4}).$$

We denote by p the projection

$$p : S(t_4) \rightarrow \tilde{t}_3^{1,3,4} \otimes S(t_3^{2,3,4}),$$

which is the tensor product of: the identity on the last factor, the projection to degree 1 on the factor $S(\tilde{t}_3^{1,3,4})$, and the projection to degree 0 in all other factors. We also denote by $m : \tilde{t}_3^{1,3,4} \otimes S(t_3^{2,3,4}) \rightarrow S(t_3)$ the map induced by the identifications $\tilde{t}_3^{1,3,4} \subset \tilde{t}_3^{1,3,4} \simeq t_3$, $\tilde{t}_3^{2,3,4} \simeq t_3$ followed by the product map in $S(t_3)$. We denote by d_1, d_2, d_3 the maps $\mathcal{T}_3 \rightarrow \mathcal{T}_4$ defined by

$$\begin{aligned} d_1(x) &= x^{12,3,4} - x^{1,3,4} - x^{2,3,4}, \\ d_2(x) &= x^{1,23,4} - x^{1,2,4} - x^{1,3,4}, \\ d_3(x) &= x^{1,2,34} - x^{1,2,3} - x^{1,2,4}, \end{aligned}$$

so $d = d_1 - d_2 + d_3$. The maps d_i are compatible with the filtrations of $\mathcal{T}_3$ and $\mathcal{T}_4$; we denote by $\text{gr}^k(d_i)$ the corresponding graded maps, so $\text{gr}^k(d) = \text{gr}^k(d_1) - \text{gr}^k(d_2) + \text{gr}^k(d_3)$. Then if we set

$$x_1 = \sum_{a,b,c|a+b+c=k-1} (t^{1,2})^a (t^{1,3})^b (t^{2,3})^c \otimes e_{a,b,c},$$

where $e_{a,b,c} \in \tilde{t}_3$, we have

$$m \circ p \circ \text{gr}^k(d_1)(x) = \left(\sum_{\alpha=0}^k \alpha x_\alpha \right) - (t^{2,3})^{k-1} e_{0,0,k-1}.$$

On the other hand, let us define the i -degree of an element of $(\tilde{t}_{|I|})^I$ to be 1 if $i \in I$ and 0 if $i \notin I$. Then the i -degree of $\bigotimes_{I \subset \{1,\dots,n\}} S^{\alpha_I}((\tilde{t}_{|I|})^I) \subset S(t_n)$ is $\sum_{I|i \in I} \alpha_I$. If x is homogeneous

for the 1-degree, then so is $\text{gr}^k(d_2)(x)$, and $1\text{-degree}(\text{gr}^k(d_2)(x)) = 1\text{-degree}(x)$. On the other hand, the elements of $S(t_4)$ whose 1-degree is $\neq 1$ are in the kernel of p . It follows that

$$m \circ p \circ \text{gr}^k(d_2)(x_\alpha) = 0 \text{ if } \alpha \neq 1,$$

and $p \circ \text{gr}^k(d_2)(x_1) = (e_{0,0,k-1})^{1,3,4} [(t^{2,4} + t^{3,4})^{k-1} - (t^{3,4})^{k-1}]$, so

$$m \circ p \circ \text{gr}^k(d_2)(x_1) = e_{0,0,k-1} [(t^{1,3} + t^{2,3})^{k-1} - (t^{2,3})^{k-1}].$$

Finally, $p \circ \text{gr}^k(d_3)(x) = 0$. If x is such that $\text{gr}^k(d)(x) = 0$, we have $m \circ p \circ \text{gr}^k(d)(x) = 0$, so

$$\sum_{\alpha \geq 0} \alpha x_\alpha = e_{0,0,k-1} (t^{1,3} + t^{2,3})^{k-1}.$$

Looking at degrees in the decomposition (7.14), we get $x_\alpha = 0$ for $\alpha \geq 2$, and $x_1 = e_{0,0,k-1}(t^{1,3} + t^{2,3})^{k-1}$. Using the projection $p' : S(t_4) \rightarrow \tilde{t}_3^{1,2,4} \otimes S(t_3^{1,2,3})$, we get in the same way $x_1 = e_{k-1,0,0}(t^{1,2} + t^{1,3})^{k-1}$. Now $e_{k-1,0,0}(t^{1,2} + t^{1,3})^{k-1} = e_{0,0,k-1}(t^{1,3} + t^{2,3})^{k-1}$ implies $e_{k-1,0,0} = e_{0,0,k-1} = 0$ so $x_1 = 0$. Therefore $x \in S^k(\bigoplus_{1 \leq i < j \leq 3} \mathbb{K}t^{i,j})$. Let us set

$x = S(t^{1,2}, t^{1,3}, t^{2,3})$, where S is a homogeneous polynomial of degree k of $\mathbb{K}[u, v, w]$. Since $d(x) = 0$, we have

$$\begin{aligned} & S(t^{1,3} + t^{2,3}, t^{1,4} + t^{2,4}, t^{3,4}) - S(t^{1,2} + t^{1,3}, t^{1,4}, t^{2,4} + t^{3,4}) \\ & + S(t^{1,2}, t^{1,3} + t^{1,4}, t^{2,3} + t^{2,4}) = S(t^{2,3}, t^{2,4}, t^{3,4}) + S(t^{1,2}, t^{1,3}, t^{2,3}) \end{aligned}$$

(equality in $S(\bigoplus_{1 \leq i < j \leq 4} \mathbb{K}t^{i,j})$).

Applying $\frac{\partial}{\partial t^{1,2}} \circ \frac{\partial}{\partial t^{3,4}}$ to this equality, we get

$$(\partial_u \partial_w S)(t^{1,2} + t^{1,3}, t^{1,4}, t^{2,4} + t^{3,4}) = 0,$$

therefore $\partial_u \partial_w S = 0$. We have therefore

$$S(u, v, w) = P(u, v) + Q(v, w),$$

where P and Q are homogeneous polynomials of degree k . Moreover, $d(x) = 0$, so

$$\begin{aligned} & [P(t^{1,2}, t^{1,3} + t^{1,4}) - P(t^{1,2} + t^{1,3}, t^{1,4}) - P(t^{1,2}, t^{1,3})] \\ & + [Q(t^{1,4}, t^{2,4}, t^{3,4}) - Q(t^{1,4}, t^{2,4} + t^{3,4}) - Q(t^{2,4}, t^{3,4})] \\ & + [P(t^{1,3} + t^{2,3}, t^{1,4} + t^{2,4}) + Q(t^{1,3} + t^{1,4}, t^{2,3} + t^{2,4}) - P(t^{2,4}, t^{2,4}) - Q(t^{1,3}, t^{2,3})] = 0. \end{aligned} \quad (7.15)$$

Write this as an identity

$$B(t^{1,2}, t^{1,3}, t^{1,4}) + C(t^{1,4}, t^{2,4}, t^{3,4}) + A(t^{2,3}, t^{1,4}, t^{1,3}, t^{2,4}) = 0.$$

Then A (resp., B, C) is independent on $t^{2,3}$ (resp., $t^{1,2}, t^{3,4}$). Let us now determine P and Q . Since $B(t^{1,2}, t^{1,3}, t^{1,4}) = B(0, t^{1,3}, t^{1,4})$, we have $P(u, v + w) - P(u + v, w) - P(u, v) = P(0, v + w) - P(v, w) - P(0, v)$. Therefore $(d\bar{P})(u, v, w) = 0$, where $\bar{P}(u, v) = P(u, v) - P(0, v)$ and d is the co-Hochschild differential of polynomials in one variable. The corresponding cohomology is zero, so we have a polynomial $\bar{P}$, such that

$$P(u, v) - P(0, v) = \bar{P}(u + v) - \bar{P}(u) - \bar{P}(v).$$

We conclude that $P(u, v)$ has the form

$$P(u, v) = \bar{P}(u + v) - \bar{P}(u) - R(v) \quad (7.16)$$

where $\bar{P}$ and R are polynomials in one variable of degree k ; since $P(u, v)$ is homogeneous of degree k , we can assume that $\bar{P}$ and R are monomials of degree k . In the same way, since $C(t^{1,4}, t^{2,4}, t^{3,4}) = C(t^{1,4}, t^{2,4}, 0)$, we have $Q(u + v, w) - Q(u, v + w) - Q(v, w) = Q(u + v, 0) - Q(u, v) - Q(v, 0)$, so $(d\tilde{Q})(u, v, w) = 0$, where $\tilde{Q}(u, v) = Q(u, v) - Q(u, 0)$. So $Q(u, v)$ has the form

$$Q(u, v) = \tilde{Q}(u + v) - \tilde{Q}(v) - S(u), \quad (7.17)$$

where $\bar{Q}$ and S are polynomials in one variable of degree k , which can be assumed to be monomials of degree k . We have therefore

$$x = \bar{P}^{1,23} + \bar{Q}^{12,3} - \bar{P}^{1,2} - \bar{Q}^{2,3} - T^{1,3},$$

where $\bar{P} = \bar{P}(t^{1,2})$, $\bar{Q} = \bar{Q}(t^{1,2})$ and $T = (R + S)(t^{1,2})$. So $x = d(\bar{Q}) + (\bar{P} + \bar{Q})^{1,23} - (\bar{P} + \bar{Q})^{1,2} - T^{1,3}$. Set $a = \bar{P} + \bar{Q}$; we have $d(y) = 0$, where $y = a^{1,23} - a^{1,2} - T^{1,3}$; applying ε_1 to $d(y) = 0$, we get $T^{2,3} - T^{2,4} = 0$, so $T = 0$. We then get $a^{12,34} - a^{12,3} - a^{2,34} + a^{2,3} = 0$. Applying $\varepsilon_3 \circ \varepsilon_2$ to this identity, we get $a^{1,4} = 0$. Finally $\bar{P} = -\bar{Q}$, so $x = d(\bar{Q})$, which proves the lemma. $\square$

- c - Isotropy groups

Proposition 7.3 can be generalized to the case of a pair of elements of Pent_{Lie} , and it implies that the isotropy group of each element of Pent_{Lie} is the additive group $\{e^{\lambda t^{1,2}}, \lambda \in \mathbb{K}\}$. Let Φ be an element of Pent . There exists an element Φ_{Lie} of Pent_{Lie} in the orbit of Φ . So the isotropy groups of Φ and Φ_{Lie} are conjugated. Since $1 + (\mathcal{T}_2)_0$ is commutative, the isotropy group of Φ is $\{e^{\lambda t^{1,2}}, \lambda \in \mathbb{K}\}$. $\square$

ACKNOWLEDGEMENTS

B.E. would like to thank Y. Kosmann-Schwarzbach for communicating to him [Dr4], where the question of the construction of quasi-groups is raised, as well as for discussions on this question in 1995. Both authors would also like to thank P. Etingof, F. Gavarini, Y. Kosmann-Schwarzbach and P. Xu for discussions.

References

- [Ban] Bangoura M. *Quasi-groupes de Lie-Poisson*. C.R. Acad. Sci. Paris Sér. I Math. **319** (1994), no. 9, 975–978.
- [Dr1] Drinfeld V. “Quantum groups”. *Proceedings of the International Congress of Mathematicians*, Amer. Math. Soc., Providence, RI **1, 2** (1987), 798–820.
- [Dr2] Drinfeld V. “Quasi-Hopf algebras”. *Leningrad Math. J.* **1** (1990), no. 6, 1419–1457.
- [Dr3] Drinfeld V. “On quasitriangular quasi-Hopf algebras and on a group closely connected with $\text{Gal}(\bar{\mathbb{Q}}/\mathbb{Q})$ ”. *Leningrad Math. J.* **2** (1991), no. 4, 829–860.
- [Dr4] Drinfeld V. letter to Y. Kosmann-Schwarzbach, 1991.
- [EGH] Enriquez B., Gavarini F., Halbout G. “On the unicity of braidings of quasitriangular Lie bialgebras”, prépublication de l’IRMA.
- [EH] Enriquez B., Halbout G. “An $\hbar$ -adic valuation property of universal R -matrices”, to appear in *J. of Algebra*.
- [Ga] Gavarini F. “The quantum duality principle”, *Ann. Inst. Fourier* **52** (2002), no. 3, 1–26.
- [GH] Gavarini F., Halbout G. “Tressages des groupes de Poisson formels à dual quasitriangulaire”, *J. Pure Appl. Algebra* **161** (2001), no. 3, 295–307.

- [KS] Kosmann-Schwarzbach, Y. “Poisson-Lie groups and beyond”, *J. Math. Sci.* **82** (1996), no. 6, 3807–3813.
- [WX] Weinstein A., Xu P. “Classical Solutions of the Quantum Yang-Baxter Equation”, *Commun. Math. Phys.* 148 (1992), 309–343.